

Minimally Invasive Plate Osteosynthesis: A Review

Pradeep Choudhari^{1*}, Mudit Baxi²

¹Professor & Unit Head, ²PG Student, Dept. of Orthopaedics, Sri Aurobindo Medical College & PG Institute, Indore

*Corresponding Author:

Email: pchoudhari@rediffmail.com

Abstract

Present-day concepts of osteosynthesis emphasize upon biological healing of fractures. This has led to advent of minimally invasive plate osteosynthesis. This technique if used when indicated leads to numerous advantages such as higher rates of union with decreased surgical time, blood loss, post-operative pain, infection rates and certain other postoperative complications. This technique is being applied commonly in fractures such as shaft of humerus, proximal and distal femur, proximal and distal tibia with several benefits over its conventional counterparts. In this review, the authors highlight upon historical aspect of development of this technique, its implementation in different fractures, various factors determining its stable application and its pearls and pitfalls.

Access this article online	
Quick Response Code:	Website: www.innovativepublication.com
	DOI: 10.5958/2395-1362.2016.00029.3

Introduction

History of plating dates back to 1886 when the first reported use of a plate for internal fixation of fracture was accomplished by Hansman^[1]. The science of plating flourished for several years till the early twenties following an era of disfavor. Plating returned to vogue following the introduction of revolutionary concepts by Muller. He and his colleagues, under the banner of Arbeitsgemeinschaft für Osteosynthese (AO) foundation laid down the concepts of operative fixation of fractures. Complete restoration of the bony anatomy under direct vision during fracture fixation was the rule. This resulted in direct fracture healing with absolute stability^[2]. But, this came at a price. The price was the risk of bone or soft tissue necrosis and delayed healing as there was widespread soft tissue dissection and interruption of periosteal blood flow^[3].

This led to advent to more biology friendly techniques. These biological techniques lay stress upon maximal preservation of blood supply around the fractured bone by minimal direct handling of the fracture environment^[4-5]. They offer advantages which include preservation of the fracture hematoma, less surgical trauma to the surrounding soft tissues^[6], reduced operative time therefore decreased risk of infection and ultimately leading to indirect bone healing with abundant callus.

Minimally invasive plate osteosynthesis (MIPO) is a novel technique for application of principles of biological fracture healing with a philosophy of dealing with soft tissues with utmost respect. Brunner and

Weber are credited for popularizing this method in the early eighties.^[7] Further later, following advancements in implant technology and development of techniques of indirect fracture reduction this technique gained more popularity.

The evolution of MIPO began with the use of bridge plating^[8]. In initial days a conventional plate was applied for comminuted femoral fractures using long incisions but with preservation of the vastus lateralis muscle resulting in healing with abundant callus^[9]. Subsequently, the size of incisions got decreased, rather two small incisions were given at proximal and distal ends and plates were bridged sub muscularly over the fracture fragments^[10-12].

This technique causes minimal distress to soft tissues and bone, provides access to the bone through soft tissue windows, with minimal or no contact with the fracture by indirect reduction tools and leaving behind minimal foot prints.

The study focuses over various aspects of application of MIPO technique used in commonly indicated fractures, along with factors influencing its mechanical stability, recent advances and its downside.

What are the indications of MIPO?

Although minimally invasive approach for plating bears numerous advantages, due to its limitations it is not universally applicable. A careful preoperative planning is required prior to its implementation with bearing in mind the anatomical aspect of the fracture, condition of soft tissue, etiology of the fracture, type of construct required and date since injury. Indications for MIPO have been summarized by Babst and Khong^[13]:

- epi-/metaphyseal fractures;
- poor soft tissue condition obviating an open procedure;
- conditions where intramedullary nailing is not feasible: unsuitable fracture pattern (intra-articular extension), narrow, deformed or obstructed intra-medullary canal;

- cases in which intramedullary reaming is contraindicated or can be detrimental (polytrauma patients with respiratory compromise);
- fractured bones with other implants or prosthesis in situ;
- Settings where of an image intensifier is not available.

Regional Considerations

Shaft Humerus: A wide variety of treatment modalities are available for treatment of shaft humerus: functional cast bracing^[14], open compression plating^[15], intra-medullary nailing.^[16] All these methods have shown good results when used judiciously with their respective indications.

Non operative treatment is economical and technically less demanding. But it leads to prolonged immobilisation and increased malunion rates. Open compression plating promises better reduction of fracture fragments^[17] but causes extensive soft-tissue handling, loss of fracture hematoma and thus loss of biological environment for fracture to heal^[18]. It is also associated with complications such as sepsis, iatrogenic radial nerve injury^[19] and nonunion. It is also a technically demanding procedure requiring long learning curve^[20].

Use of MIPO by the means of anterior bridge plating helps in preservation of biological atmosphere and thus promotes adequate healing at the fracture site. Surgery is minimally invasive so there is less blood loss, neurovascular complications, post-operative pain, chances of sepsis and lesser hospital stay. In the middle third fractures, MIPO is safer from the risks of iatrogenic radial nerve injuries. However, it is not the case with distal third fractures as there is risk of radial nerve entrapment under the plate^[21]. Since exposure is minimal chances of infection encountered are less too.

Proximal Femur: The functional outcomes following plating of subtrochanteric fractures are far more inferior to those following the intramedullary devices^[22-23]. But nailing in certain complex fracture can be technically demanding. In such situations plating can be beneficial. Additionally plates are less expansive as compared to nails. Other relative indications are cases with small medullary canals or cases of poly trauma involving the pulmonary system in which reaming can be detrimental^[24]. In such situations plating can be done with minimal access to have the biological advantages. Implants such as DCS, Condylar buttress plate can be used with satisfactory outcome with MIPO technique^[25].

Distal Femur: The conventional lateral approach to distal femur offers the window for excellent visualization and direct reduction of the fracture. But this comes at a price of massive soft tissue exposure, loss of blood, devitalisation of fracture fragments and

need for bone grafting in certain circumstances ultimately leading to infections, non-unions^[26].

But with the advent of various minimally invasive procedures in different forms has led to decreased incidence of such complications. Various techniques have been devices such as MIPO, MIPPO, TARPO, LISS offering minimal of complications.

MIPO is convenient for cases with extra articular involvement with complex Meta/diaphyseal fragments. MIPPO (minimally invasive percutaneous plate osteosynthesis) technique encompasses use of specialized instruments for application of implants such as DCS. Thus, offering advantage of DCS with minimal soft tissue handling^[10].

TARPO (Transarticular Approach and Retrograde Plate Osteosynthesis) offers direct visualization of articular surface with minimal access to the metaphyseal region. Hence, its advantage lies in osteosynthesis of articular fractures. It utilizes the use of an incision given for lateral parapatellar arthrotomy for direct visualization for restoration of the articular congruity and an indirect fixation of the diaphyseal component^[27]. Although these techniques are advantageous, but they are technically more demanding.

LISS for distal femur (Less invasive stabilization system) comprises of the use of an anatomically contoured plate which is inserted via minimal access into the sub-muscular plane through an aiming device after indirect reduction of the fracture^[28]. Screws are inserted using the aiming device with minimal incisions. Thus the advantages of minimal tissue handling are offered, but the technique is costly and requires availability of high profile instrumentation.

Proximal Tibia: High velocity tibial plateau fractures often encompass a combination of complex articular involvement with substantial comminution and associated severe soft tissue injuries. Operating upon such injuries with conventional approaches in presence of the damaged soft tissues leads to incidence of high amount complications such as infections^[29], wound dehiscence, hardware prominence. To avoid such situations a two staged approach, use of dual incisions or definitive external fixation is preferred. These methods can increase morbidity and are not cost effective as multiple surgeries are required and increased hospital stay is required.

The usefulness of minimally invasive approaches comes to advantage in such situations. Here, MIPO has shown lower incidence of soft-tissue complications and yields in better end-results than ORIF^[30-31].

Distal Tibia: Plating is gold standard in operative treatment of distal tibial fractures. Intramedullary nailing is not preferred treatment for distal tibial fractures as these fractures commonly extend to the tibial pilon^[32]. Application of external fixators in distal

tibial fractures is commonly associated with a high complication rate. Complications such as pin tract infection pin loosening and high rate of malunion are common^[33].

Optimal treatment of distal tibial fractures is challenging owing to its fragile soft tissue cover and tenuous blood supply. Cadaveric studies have shown that its vascularity is derived from extra osseous anastomosis of branches of anterior and posterior tibial arteries. These arteries arrive into distal tibia from the medial side. Open plating can thus lead to disruption of this blood supply and hamper the vascularity^[34].

MIPO has shown to have higher of union rates and lower complication rates as compared with the conventional open method^[35-36].

Factors Influencing Stability of Construct^[37]

In order to achieve a stable construct the following guidelines are recommended:

Plate Length: Plate length should be two to three times bigger than length of the fracture in comminuted fractures and eight to ten times higher in simple fractures. Use of such longer constructs is requirements for an ideally stable construct. The length increases the pullout force acting on the screw because of long lever arm for each screw. MIPO with its inherent ability of plate insertion with minimal incisions offers such advantage of placement of larger plates without extensive exposure.

Number of screws: Two screws on each side of the fractures are prerequisite for a stable construct. But, being on the safer side it is recommend to use three screws on either side so as to take care of chances of failure due to screw breakage. A plate screw density below 0.4 to 0.5 is recommended, this implies that less than half of the plate holes are occupied by screws

Locking versus non locking screws: Use of locking screws is recommended especially in bones with poor quality as locking heads provide strength against not only pullout but also bending.

Bicortical versus unicortical screws: A purchase of certain amounts of threads in the engaging cortex is necessary for firm anchorage of plate to the bone surface. Healthy cortices are dense enough for a good purchase. In such bones even a monocortical screw can resist the torsional forces acting over the fracture site. Whereas, in case of porous bones the cortex is thin. Here, even minimal torsional forces can lead to instability as the screw pullout strength becomes lower due to decreased working length of the screw. Therefore, in such cases use of bicortical screw is recommended.

Self-drilling versus non self-drilling screws: Use of self-drilling screws is recommended only for unicortical screws as they can cause soft issue damage while drilling far cortex for bicortical applications.

Screw direction: Parallel insertion has lesser strength against pull-out, as a pull-out force is applied, entire screws and plate interface pulls out as a whole. The pull-out strength can be increased by use of screw in diverging directions^[38].

Pearls for performing MIPO

1. In order to achieve desirable outcome a few steps are critical: performing *MIPO earlier the better* as reduction and achievement of adequate length becomes difficult by indirect measures in older fractures; having *accurate knowledge of surface anatomy* as precise incisions and dissection is required; achievement of indirect closed reduction, *avoiding exposure of the fracture* and using *special tools* if required (described below); making *small incisions* for implant insertion; use of plates preferably *anatomically contoured* with *low bone contact* as pure *splints*; *elastic bridging* of the fracture to formulate a relatively stable construct^[39].
2. Attainment of intra-operative control over bone alignment and rotation of fragments can be cumbersome even to the experienced hands. Repeated use of intra-operative use of fluoroscopy exposes the surgeon to hazards of radiation^[13]. The following few signs can minimize the hassle while using fluoroscopy- **Cortical step sign:** the opposite bone ends have unlike cortical thickness; **Diameter step sign:** in presence of malrotation, diameters of proximal and distal ends appear to be of different dimensions; **lesser trochanter sign:** limb rotation leads to altered appearance of the lesser trochanter^[40].
3. In order to minimize these hardships, development of better instruments and techniques is essential. These include better navigation tools, plates with newer types of instrumentation such as aiming devices, indirect reduction tools such as clamps, pointed forceps such as collinear forceps, use of distracters, wire passers, modified t handles and the f-tool.^[41-42]
4. Location of a plate hole without Fluoroscopy: Fluoroscopy is not required to locate a plate hole. Place identical plate over the subcutaneous plate and locate holes, then make small stab incisions. Use small hemostat to locate the holes.

Pitfalls

1. As the fracture site is not directly visualized, chances of mal-reduction exist which lead to malunion of fractures^[43]. Also, use of this

- technique limits its application in articular fractures where direct visualization of the joint surface is required. In such cases this technique is used in conjugation with standard open approaches.
2. As the procedure involves meticulous soft tissue handling and minimal exposure the procedure is considered technically demanding and a longer learning curve is required^[44-45].
 3. This technique is contraindicated in cases of open fractures with severe soft tissue loss with no coverage of exposed bone, as the fracture site is already exposed, associated neuro-vascular injuries as exploration of injured structure is required, pathological fractures and cases of non-union and delayed unions as these fractures require freshening of fracture margins with or without bone grafting.
 4. As compared to conventional plating this procedure requires higher fluoroscopic control and thus exposes the surgeon to higher amount of radiation^[13].
 5. Implant removal can be tedious, as small incisions are required for plate insertion and often incisions are not in place of position of screws as swelling subsides. Large extensile incisions are then required for implant removal^[46].

Conclusion

MIPO is simple, safe and effective method of treatment for diaphyseal & metaphyseal fractures. It does not require direct visualization of the fracture at the cost of extensive soft tissue dissection. We are confident in suggesting this alternative approach of plating.

References

1. Schatzker J. Osteosynthesis in trauma. *International orthopaedics*. 1996 Jul 1;20(4):244-52.
2. Perren SM. Fracture healing. The evolution of our understanding. *Acta chirurgiae orthopaedicae et traumatologiae Cechoslovaca*. 2008 Aug;75(4):241-6.
3. An Z, Zeng B, He X, Chen Q, Hu S. Plating osteosynthesis of mid-distal humeral shaft fractures: minimally invasive versus conventional open reduction technique. *International orthopaedics*. 2010 Jan 1;34(1):131-5.
4. Weller S. Biological fracture fixation-What is this. Is it another traumatological fashion or an important aspect of operating technique. 1997.
5. Xu H, Xue Z, Ding H, Qin H, An Z. Callus Formation and Mineralization after Fracture with Different Fixation Techniques: Minimally Invasive Plate Osteosynthesis versus Open Reduction Internal Fixation. *PloS one*. 2015 Oct 7;10(10):e0140037.
6. Wagner M, Frenk A, Frigg R. Locked plating: Biomechanics and biology and locked plating: Clinical indications. *Techniques in Orthopaedics*. 2007 Dec 1;22(4):209-18.
7. Brunner CF, Weber BG, Brunner CF, Weber BG. Antigleitplatte. Besondere Osteosynthesetechniken. 1981:115-33.
8. Heitemeyer U, Kemper F, Hierholzer G, Haines J. Severely comminuted femoral shaft fractures: treatment by bridging-plate osteosynthesis. *Archives of orthopaedic and traumatic surgery*. 1987 Aug 1;106(5):327-30.
9. Bolhofner BR, Carmen B, Clifford P. The results of open reduction and internal fixation of distal femur fractures using a biologic (indirect) reduction technique. *Journal of orthopaedic trauma*. 1996 Aug 1;10(6):372-7.
10. Krettek C, Schandelmaier P, Nliclau T, Tscherne H. Minimally invasive percutaneous plate osteosynthesis (MIPPO) using the DCS in proximal and distal femoral fractures. *Injury*. 1997 Dec 31;28:A20-30.
11. Wenda K, Runkel M, Degreif J, Rudig L. Minimally invasive plate fixation in femoral shaft fractures. *Injury*. 1997 Dec 31;28:A13-9.
12. Baumgaertel F, Buhl M, Rahn BA. Fracture healing in biological plate osteosynthesis. *Injury*. 1998 Dec 31;29:3-6.
13. Babst R, Khong K. Minimally invasive surgery. *AO Principles of Fracture Management*. Stuttgart, Germany. 2007:199-212.
14. Camden P, Nade S. Fracture bracing the humerus. *Injury*. 1992 Jan 1;23(4):245-8.
15. Bell MJ, Beauchamp CG, Kellam JK, Mc Murtry RY. The results of plating humeral shaft fractures in patients with multiple injuries. The Sunnybrook experience. *Journal of Bone & Joint Surgery, British Volume*. 1985 Mar 1;67(2):293-6.
16. Flinkkilä T, Hyvönen P, Lakovaara M, Linden T, Ristiniemi J, Hämäläinen M. Intramedullary nailing of humeral shaft fractures: a retrospective study of 126 cases. *Acta orthopaedica Scandinavica*. 1999 Jan 1;70(2):133-6.
17. Zhiquan A, Bingfang Z, Yeming W, Chi Z, Peiyan H. Minimally invasive plating osteosynthesis (MIPO) of middle and distal third humeral shaft fractures. *J Orthop Trauma* 2007;21:628-33
18. Pospula W, Abu Noor T. Percutaneous fixation of comminuted fractures of the humerus: initial experience at Al Razi hospital, Kuwait. *Med Princ Pract* 2006;15:423-6.
19. Paris H, Tropiano P, Clouet D'orval B, Chaudet H, Poitout DG. Fractures of the shaft of the humerus: systematic plate fixation. Anatomic and functional results in 156 cases and a review of the literature. *Rev Chir Orthop Reparatrice Appar Mot*. 2000 Jun;86(4):346-59.
20. Kumar C, Gupta AK, Nath R, Ahmad J. Open reduction and locking plate fixation of displaced proximal humerus fractures. *Indian journal of orthopaedics*. 2013 Mar 1;47(2):156.
21. Malhan S, Thomas S, Srivastav S, Agarwal S, Mittal V, Nadkarni B, Gulati D. Minimally invasive plate osteosynthesis using a locking compression plate for diaphyseal humeral fractures. *Journal of Orthopaedic Surgery*. 2012;20(3).
22. Brien WW, Wiss DA, Becker V Jr, et al. Subtrochanteric femur fractures: A comparison of the Zickel nail, 95 degrees blade plate, and interlocking nail. *J Orthop Trauma*. 1991;5(4):458-464.
23. Forward DP, Doro CJ, O'Toole RV, et al. A biomechanical comparison of a locking plate, a nail, and a 95° angled blade plate for fixation of subtrochanteric femoral fractures. *J Orthop Trauma*. 2012;26(6):334-340.
24. Prasarn ML, Ahn J, Helfet DL et al. Bisphosphonate-associated femur fractures have high complication rates with operative fixation. *Clin Orthop Relat Res*. 2012;470(8):2295-2301.

25. Sirbu PD, Carata E, Petreus T, Munteanu F, Popescu C, Asaftei R, Botez P. Dynamic Condylar Screw (DCS) versus Condylar Blade Plate (CBP) in Complex Supracondylar Femoral Fractures—A Biomechanical Study. In International Conference on Advancements of Medicine and Health Care through Technology 2009 (pp. 409-412). Springer Berlin.
26. Ostrum RF, Geel C. Indirect reduction and internal fixation of supracondylar femur fractures without bone graft. *Journal of orthopaedic trauma*. 1995 Aug 1;9(4):278-84.
27. Krettek C, Miclau T, Stephan C, Tscheme H. Transarticular Approach and Retrograde Plate Osteosynthesis (TARPO): An Alternative Surgical Approach for Complex Distal Intra-Articular Femur Fractures. *Techniques in Orthopaedics*. 1999 Sep 1;14(3):219-29.
28. Schuetz M, Müller M, Krettek C, Höntzsch D, Regazzoni P, Ganz R, Haas N. Minimally invasive fracture stabilization of distal femoral fractures with the LISS: A prospective multicenter study results of a clinical study with special emphasis on difficult cases. *Injury*. 2001 Dec 31;32:48-54.
29. Mallik AR, Covall DJ, Whitelaw GP. Internal versus external fixation of bicondylar tibial plateau fractures. *Orthopaedic review*. 1992 Dec;21(12):1433-6.
30. Stannard JP, Wilson TC, Volgas DA, Alonso JE. The less invasive stabilization system in the treatment of complex fractures of the tibial plateau: short-term results. *J Orthop Trauma* 2004;18:552-8.
31. Cole PA, Zlowodzki M, Kregor PJ. Treatment of proximal tibia fractures using the less invasive stabilization system: surgical experience and early clinical results in 77 fractures. *J Orthop Trauma* 2004;18:528-35.
32. Krackhardt T, Dilger J, Flesch I, Höntzsch D, Eingartner C, Weise K. Fractures of the distal tibia treated with closed reduction and minimally invasive plating. *Arch Orthop Trauma Surg* 2005;125:87-94.
33. Siebenrock KA, Schillig B, Jakob RP. Treatment of complex tibial shaft fractures. Arguments for early secondary intramedullary nailing. *Clin Orthop Relat Res* 1993;290:269-74.
34. Borrelli J Jr, Prickett W, Song E, Becker D, Ricci W. Extra osseous blood supply of the tibia and the effects of different plating techniques: a human cadaveric study. *J Orthop Trauma* 2002;16(10):691-5.
35. Oh CW, Kyung HS, Park IH, Kim PT, Ihn JC. Distal tibia metaphyseal fractures treated by percutaneous plate osteosynthesis. *Clin Orthop* 2003;408:286-91
36. Helfet DL, Suk M. Minimally invasive percutaneous plate osteosynthesis of fractures of the distal tibia. *Instructional course lectures*. 2003 Dec;53:471-5.
37. Gautier E, Sommer C. Guidelines for the clinical application of the LCP. *Injury*. 2003 Nov 30;34:63-76.
38. Cronier P, Pietu G, Dujardin C, Bigorre N, Ducellier F, Gerard R. The concept of locking plates. *Orthopaedics & traumatology: surgery & research*. 2010 Jun 30;96(4):S17-36.
39. Thakur AJ. *Elements of Fracture Fixation*. Elsevier Health Sciences; 2015 Nov 10.
40. Krettek C, Miclau T, Gru O, Schandelmaier P, Tscherne H. Intraoperative control of axes, rotation and length in femoral and tibial fractures technical note. *Injury*. 1998 Dec 31;29:29-39.
41. Apivatthakakul T, Phornphutkul C, Patumasutra S. Idea and innovation: Simple minimally invasive plate osteosynthesis (MIPO) instruments. *Injury Extra*. 2009 Feb 28;40(2):39-44.
42. Gupta RK, Rohilla RK, Sangwan K, Singh V, Walia S. Locking plate fixation in distal metaphyseal tibial fractures: series of 79 patients. 2010;34(8):1285-1290.
43. Shin SJ, Sohn HS, Do NH. Minimally invasive plate osteosynthesis of humeral shaft fractures: a technique to aid fracture reduction and minimize complications. *Journal of orthopaedic trauma*. 2012 Oct 1;26(10):585-9.
44. Anuar-Ramdhan IM, Azahari IM. Minimally Invasive Plate Osteosynthesis with Conventional Compression Plate for Diaphyseal Tibia Fracture. *Malaysian orthopaedic journal*. 2014 Nov;8(3):33.
45. Francois J, Vandeputte G, Verheyden F, Nelen G. Percutaneous plate fixation of fractures of the distal tibia. *Acta orthop belg*. 2004;70(2):148-154.
46. Georgiadis GM, Gove NK, Smith AD, et al (2004) Removal of the less invasive stabilization system. *J Orthop Trauma*; 18(8):562-564.