

Prevalence of anencephaly associated anomalies

Ganesh T Waghmode^{1,*}, Vishal M. Salve², Anjali G. Gosavi³

¹Associate Professor, ^{2,3}Professor, Dept. of Anatomy, ^{1,2}Navodaya Medical College, Raichur, Karnataka ³Ashwini Rural Medical College, Solapur, India

***Corresponding Author:**

Email: gw1726@gmail.com

Abstract

Introduction: Defective closure of rostral pore of neural tube leads to anencephaly. Folic acid deficiency is most common cause anencephaly. In most of the cases of Anencephaly is associated with other systemic anomalies.

Objectives: Aim of present study was to find out the incidence of anencephaly associated systemic anomalies and their correlation with maternal age, sex of the foetus.

Materials and Methods: 32 anencephaly fetuses obtained from the Department of obstetrics and gynaecology Navodaya Medical College. The period of study was from January 2013 to December 2016.

Results: Out of 32 cases, 25(78.12%) cases showed presence of systemic anomalies associated with anencephaly, 53.12% of the cases were observed in primigravida. Most common associated anomaly was spinabifida 28.12% followed by gastrointestinal anomalies 6.25%.

Conclusion: Incidence of anencephaly associated anomalies is not uncommon; hence gross systemic examination of the anencephalic abortus is required to find out incidence of anencephaly associated anomalies.

Keywords: Anencephaly, Craniospinal rachinoschisis, Exomphalos, Spina bifida.

Received: 12th July, 2017

Accepted: 7th September, 2017

Introduction

Anencephaly is a neural tube defect which is due to the defective closure of cranial neuropore. Closure of the cranial neuropore takes place between the third and fourth week of gestation at 18-20 somite stage.¹ Anencephaly is most common congenital malformation during foetal development. The prevalence of anencephaly in India is 1 to 5 per 1000 live births.²

Present study was undertaken to determine the prevalence of anencephaly among congenital malformations. Organs from infants with anencephaly can be used for organ transplantation. David A et al have discussed anencephaly infant and their use in organ transplant, but most frequently anencephaly is associated with other anomalies. In present study we have correlated anencephaly with associated systemic anomalies, maternal age, birth order and sex of the foetus. Our study will be helpful in selecting anencephaly infant for organ transplantation.³

The Anencephaly shows a heterogeneous aetiology, ranging from environmental to genetic causes. Genetic disorders related with folic acid metabolism, glucose metabolism and inositol metabolism are associated with neural tube defects.⁴

Aetiology of Meroanencephaly, holoanencephaly, craniorachischisis and spinabifida was described by multisite closure model of neural tube of Van Allen et al. Anencephaly is commonly associated with other systemic anomalies; only few studies are available on anencephaly associated disorders in India.⁵

Materials and Methods

32 fetuses with neural tube defect were obtained from department of obstetrics and gynaecology. This study was done for three years from January 2013 to February 2016.

During this period, the fetuses which were still born with neural defect were collected after getting the consent of the parents. Fetuses died or terminated due to other causes were excluded from the study. They were embalmed and stored in 10% formalin solution.

The fetuses were observed in detail externally for the sex, type of neural defect and other associated anomalies and the findings were documented. The approximate crown rump length was measured and the probable gestational age of the fetuses was calculated. The fetuses were dissected to find other internal anomalies. The statistical analysis was performed by using frequency and percentage.

Results

Table 1: Incidence of anencephaly and its correlation with maternal age, parity, gestational age and sex of the foetus.

| Age of mother | Parity | | Gestational age | Sex of foetus | |
|---------------|----------------------|--------------|---------------------|---------------|-----------------------|
| | 20 years to 30 years | Primigravida | | 17 (53.12%) | 20 weeks to full term |
| Gravid two | | 11 (34.37%) | Female | 23 (71.87%) | |
| Gravid three | | 3 (9.37%) | Ambiguous genitalia | 2(6.23%) | |
| Gravid four | | 1 (3.12%) | | | |

Table 2: Number of cases of anencephaly associated systemic anomalies

| S. No | System | Associated anomalies | No of cases |
|-------|-------------------------|-----------------------------|-------------|
| 1 | Gastrointestinal system | Gastroschisis | 01 (3.12%) |
| | | Exomphalos | 01(3.12%) |
| 2 | Respiratory system | Cyst in lung | 01(3.12%) |
| 3 | Musculoskeletal system | Achondroplasia | 01(3.12%) |
| | | Absence vertebral column | 01(3.12%) |
| 4 | Cardiovascular system | Single umbilical artery | 01(3.12%) |
| 5 | Nervous system | Meroanencephalic | 04(12.5%) |
| | | Holoanencephalic | 03 (9.37%) |
| | | Spinabifida | 09(28.12%) |
| | | Craniospinal rachinoschisis | 03 (9.37%) |
| | | Myelocele | 05(15.62%) |
| 6 | Head face neck | Cleft palate | 01(3.12%) |



Fig. 1: Encephalocele



Fig. 2: Craniospinal rachischisis



Fig. 3: Gastroschisis

Discussion

The prevalence of anencephaly shows great variation in time and space. Among the 194 World Health Organization member states, the percent reporting within each region is as follows: African (8/47;17%), Eastern Mediterranean (12/21;57%), European (26/53;49%), Americas (15/35;43%), South-East Asian (4/11;36%) and Western Pacific (9/27;33%).⁶

According to Mahadevan et al the prevalence of anencephaly in India is 1.8 to 7 per 1000 live births.⁷ In present study maternal age studied was between 20 years to 30 years as we have not observed anencephaly foetus with maternal age above 30 years. In a study by Isabela, et al there was no association between maternal age and anencephaly.⁸ A study by Dwight et al there was declining risk in young mother and older mothers, they observed a 'U' shape pattern while correlating anencephaly and maternal age.⁹ Vieira et al observed higher risk of spina bifida than anencephaly in younger and older mothers. Thus more detailed studies are required to correlate maternal age and anencephaly.¹⁰

In present study anencephaly associated disorders were more common in primigravida (53.12%). In a study by Vare et al Anencephaly were observed more common in primigravida.¹¹

Prevalence of anencephaly in female foetus (71.87%) was more than in male. Isabela et al observed 64% of female anencephalic foetuses in their study (y) The female preponderance also reported by Jaquier et al¹² and James et al,¹³ but study by Obeidi et al did not observed female preponderance.¹⁴

Anencephaly associated anomalies ranges from 9.4 to 43%.^{11, 16-18, 20} In present study it was higher (78.12%) than previous studies but similar to study by C. Panduranga et al 73%.¹⁵

The gastrointestinal anomalies associated with the anencephaly in present study we have observed one gastroschisis and one omphalocele (6.25%). In previous studies Golipure et al 5.3%¹⁶ and David T et al 5.7%¹⁷ gastrointestinal anomalies were similar to present study, but Vare et al mentioned (32%)¹¹ Gastrointestinal anomalies.

Respiratory anomalies in present study and study by Tan et al¹⁸ were same (3%), but majority of previous studies have not mentioned about respiratory anomalies associated with anencephaly.

Musculoskeletal disorders in present study we have observed two cases (6.25%) achondroplasia and absence of vertebral column,

but Vare et al mentioned 27%¹¹ and according to David T et al 16.3%.¹⁷

Eslavat et al¹⁹ mentioned 33.3% cardiovascular disorders associated with neural tube defects, in present study 3.12%, Tan et al 3%¹⁸ and Nielson et al 4.75%²⁰ have mentioned observations similar to present study.

Central nervous system disorders like spinabifida (28.12%), Craniospinal rachinoschisis (9.37%) and Myelocele (15.62%) were observed in present study, in previous studies only C. Panduranga et al mentioned Central nervous system disorders spinabifida (26.82%), Craniospinal rachinoschisis (12.19%) were close to present study.

3.12% of cleft palate was present in present study; previous studies mentioned similar observation Golipure et al 3.5%¹⁶ and David et al 3.5%,¹⁷ except Nielsen et al mentioned 14%²⁰ of head neck disorders.

Disorders like Diaphragmatic hernia and Genital defect were mentioned by very few previous studies; Nielsen et al 2.3%,²⁰ David et al 2.7%¹⁷ and Vare et al 5%,¹¹ in present study we have not seen these disorders.

Conclusion

Anencephaly is common in foetus of primigravid mother and most common in female foetus. Central nervous system anomalies like spinabifida (28.12%), Craniospinal rachinoschisis (9.37%) and Myelocele (5%) are commonly associated with anencephaly. Ultrasound screening in first trimester can diagnose Anencephaly and other associated systemic disorders, but macroscopic systemic examination of the abortus is required, as in most cases anencephaly is also associated with other systemic anomalies. Present study provide information of prevalence of anencephaly associated disorders, which will help in counselling of eligible couple and for management in reducing incidence of anencephaly associated disorders.

Importance of present study – Organs are required in organ transplant. Anencephalic foetuses survival is not possible, hence organs of anencephalic foetuses can be used for organ transplantation, but anencephaly is associated with other systemic anomalies. Our study will provide data of anencephaly associated disorders for counselling of eligible couple, in use of anencephalic foetuses in organ transplantation and for management in reducing incidence of anencephaly associated disorders.

Limitations of present study – We have not studied aetiology of anencephaly. We have

studied only anencephaly fetuses from hospital attached to our medical college. Duration of present study was 3years.

Conflict of Interest

No financial and other conflicting interests.

References

1. Snell. Richard. S, Lippincott Williams & Wilkis. Clinical neuroanatomy 6TH Edition 2006:511
2. JTD Hans, L Martin, H Akira. 2nd Edition. Heidelberg: Springer; 2014. Clinical Neuroembryology: Development and Developmental Disorders of the Human Central Nervous System
3. David A. S, Ronald E, Sherman, Norman C, Edwin C, Ronald P. The Infant with Anencephaly: The Medical Task Force on Anencephaly. Journal of Medicine, 1990, 322:669-74
4. Copp AJ, Greene ND, Genetics and Development of Neural tube defects. J Path 2010; 220(2):217-30
5. MI Van Allen, DK Kalousek, GF Chernoff, D Juriloff, M Harris, BC Mc Gillivray , et al. Evidence for multi-site closure of the neural tube in humans. Am J Med Genet. 1993; 47:723-43
6. World Health Organization. Global health estimates (GHE)—Disease burden. 2015. Available: http://www.who.int/healthinfo/global_burden_disease/estimates/en/index2.html. Accessed 2015 Apr 14.
7. Mahadevan B, Bhat BV. Neural tube defects in Pondicherry. Indian J Pediatr 2005; 72:557-9.
8. Isabela N, Silvia D, Ricardo B. Anencephaly: Do the Pregnancy and Maternal Characteristics Impact the Pregnancy Outcome? ISRN Obstet Gynecol. 2012 Jan; 127490
9. Dwight T. Anencephaly and maternal age, American Journal of Epidemiology, 1972, 95(4):319-26.
10. Vieira A, Castillo T. Maternal age and neural tube defects: evidence for a greater effect in spina bifida than in anencephaly. Rev Med Chil. 2005 Jan; 133(1):62-70
11. Vare AM, Bansal PC. Anencephaly. An Anatomical study of 41 anencephalies. Indian J Pediatr 1971; 38:301-5.
12. Jaquier M, Klein A, Boltshauser E. Spontaneous pregnancy outcome after prenatal diagnosis of anencephaly. *BJOG: An International Journal of Obstetrics and Gynaecology*. 2006;113(8):951-53. [PubMed]
13. James WH. The sex ratio in anencephaly. *Journal of Medical Genetics*. 1979;16(2):129-33.[PMC free article] [PubMed]
14. Obeidi N, Russell N, Higgins JR, O'Donoghue K. The natural history of anencephaly. *Prenatal Diagnosis*. 2010;30(4):357-60. [PubMed]
15. Panduranga C, Ranjit Kangle, Vijayalaxmi V. Suranagi G, Pilli S, Patil P V. Anencephaly: A pathological study of 41 cases. Journal of the Scientific Society, August 2012, Volume 39 Issue 2:81-4.
16. Gotalipoure MJ, Najafi L, Keshtkar AA. Prevalence of Anencephaly in Gorgan. Northern Iran. Arch Iran Med 2010; 13:34-7.
17. David TJ, McCrae FC, Bound JP. Congenital malformations associated with anencephaly in the Fylde peninsula of Lancashire. J Med Genet , 1983; 20:338- 41.
18. Tan KB, Tan SH, Tan KH, Yeo GS. Anencephaly in Singapore: a ten-year series 1993 – 2002. Singapore Med J 2007; 48:12-5.
19. Eslavath A, Ranga Rao, Diddi K, Chakravarthy V. Anencephaly: A 3 Years Study. Journal of Dental and Medical Sciences, Dec. 2013, Volume 12, Issue 2: 12-5
20. Nielsen LA, Maroun LL, Broholm H, Laursen H, Graem N. Neural tube defects and associated anomalies in a fetal and perinatal autopsy series. Acta Pathol Microbiol Immunol Scand 2006; 114:239-46.