

Gram positive uropathogens and their antibiogram: Data analysis at a tertiary care hospital in Karnataka

Trupti B. Naik¹, Lavanya J.^{2*}, Amruthkishan Upadhya³, Vijaykumar Mane⁴

¹Assistant Professor, Dept. of Microbiology, ChamaraJanagar Institute of Medical Sciences, Yadapura, Karnataka, ²Assistant Professor, ³Professor, Dept. of Microbiology, Subbaiah Institute of Medical College, Shivamogga, Karnataka, ⁴Assistant Professor, Dept. of Community Medicine, Koppal Institute of Medical Sciences, Koppal, Karnataka, India

***Corresponding Author:**

Email: lavanya1126@gmail.com

Abstract

Introduction: Urinary tract infections (UTI) are one of the most common infections encountered in clinical settings, being amenable to easy diagnosis with established laboratory methods. However, in recent years, antibiotic resistance has become a major public health problem worldwide owing to the extensive and inappropriate use of antimicrobial agents.

Objectives:

1. To determine the various Gram positive isolates causing urinary tract infections.
2. To determine the antibiotic susceptibility pattern of Gram positive cocci.

Material and Methods: A cross sectional study was conducted at a private tertiary care hospital in Shivamogga district of Karnataka, using secondary data of Gram positive bacteria isolated from urine samples maintained in the Microbiology laboratory registers for a period of 1 year from January 2016 to December 2016. Standard protocols were followed for bacteria isolation, identification and to assess their antibiotic susceptibility. Statistical analysis was done using MS office Excel 2010.

Results: Of the total 1254 urine specimens, 466 (37.16%) revealed positive cultures. Out of the 512 isolates from positive cultures, 196 (38.28%) were Gram positive bacteria. The predominant isolates i.e. 86 (43.87%) belonged to Enterococcus species. All the isolates were susceptible to Linezolid and Vancomycin. Majority i.e. 184 (93.87%) were sensitive to nitrofurantoin followed by gentamicin 146 (74.48%) and majority showed resistance to ampicillin i.e. 126 (64.28%).

Conclusion: The rise in the resistance among uropathogens alerts us against indiscriminate use of antibiotics, indicating the need to establish antibiotic policies along with stringent measures to ensure effectiveness of the same.

Keywords: Antibiogram, Gram positive isolates, Urinary tract infection.

Introduction

Currently, Urinary tract infections (UTI) represent one of the most commonly encountered morbidities in health care settings with an estimated 150 million cases per annum worldwide. They are caused by invasion of the urinary tract by pathogenic microorganisms.¹ The clinical manifestations depend on the part of the urinary tract affected, the causative organisms, the severity of the infection and the patient's ability to mount an immune response.² Females are more prone to UTI than males due to anatomical and physiological reasons.^{1,3,4} They occur most frequently between the ages of 16 and 35 years, with 10% of women getting an infection at some point in their lives. Recurrences are common, with nearly half of people getting a second infection within a year.⁵

Gram negative bacteria like *Escherichia coli*, Proteus species, Klebsiella species, *Pseudomonas aeruginosa*, Acinetobacter, Serratia and *Morganella morganii* are isolated from 75-95% cases of uncomplicated UTI which is most common in young, sexually active, non pregnant, premenopausal women.⁶ The remaining cases are associated with a variety of organisms, including the Gram positive bacteria like Enterococcus, Staphylococcus especially coagulase negative staphylococci, *Streptococcus agalactiae* and other less frequently isolated organisms.⁷ However,

Gram positive bacteria are found more often as etiologic agents in vulnerable groups such as pregnant women and the elderly. Complicated UTI is defined as cystitis or pyelonephritis that occurs in individuals with predisposing anatomic, metabolic or functional risk factors making their treatment difficult.⁷ and these populations are at greater risk of Gram positive and polymicrobial UTI.^{8,9}

Current management of UTIs is usually empirical, without the use of a urine culture or susceptibility testing to guide therapy. During the past decade, Gram positive cocci isolates exhibited a remarkable ability to rapidly develop antibiotic resistance and it is of great concern.⁵ Further, the Infectious Disease Society of America also recommends that physicians must keep on updating information on local susceptibility pattern of organisms causing urinary tract infections to monitor changes in their susceptibility which is a prerequisite for any hospital infection control program.^{1,3} With this background, the present study was undertaken to identify the Gram positive isolates causing UTI among patients attending a tertiary care hospital and to study their antibiotic susceptibility pattern.

Materials and Methods

A cross sectional study was conducted in the Microbiology department of a private tertiary care

hospital in Shivamogga district of Karnataka. Prior permission for the study was obtained from concerned authorities. Secondary data maintained in the laboratory registers of Microbiology department of Gram positive bacteria isolated from urine samples of patients presented with UTI for a period of 1 year from January 2016 to December 2016 were collected for the study. The following information were noted – Name, Age, Sex, Case history, Organism isolated and their antibiotic susceptibility pattern.

Statistical analysis

Analysis was done using MS Office Excel 2010.

Bacterial isolation and identification

Cases of clinically suspected urinary tract infection were routinely sent for microbiological analysis. These samples were processed in the laboratory for direct microscopy and aerobic culture and sensitivity as per the standard protocol. Routine microscopic examination of urine samples were done to look for pus cells, red blood cells and epithelial cells. Semi quantitative urine culture was done using a calibrated loop. A loopful (0.001 ml) of well mixed un-centrifuged urine was inoculated onto the surface of MacConkey agar, blood agar media and Cysteine lactose electrolyte deficient agar (CLED). All plates were then incubated at 37°C aerobically for 24 hours.

The plates were examined macroscopically for bacterial growth. Unlike Gram negative bacilli that needs more than 100 colonies corresponding to 10⁵ cfu/ml to be significant, Gram positive cocci, irrespective of the colony count were considered significant.¹⁰ The Gram positive bacterial isolates were identified using standard bacteriological tests.¹¹

Colonial appearance and morphological characters of isolated bacteria were noted and isolated colonies were subjected to preliminary tests like Gram staining and catalase test. These preliminary tests were followed by biochemical reactions for identification of the isolated organism.

Antimicrobial susceptibility testing

Antimicrobial susceptibility testing was performed on Mueller Hinton agar as per CLSI guidelines¹² and was tested for the following antibiotics - Ampicillin (10µg), Tetracycline (30µg), Ciprofloxacin (5µg), Norfloxacin (10µg), Nitrofurantoin (300µg), Gentamycin (10µg), High level Gentamycin (120µg), Linezolid (30µg), Vancomycin (30µg).

Results

Of the total 1254 urine specimens collected and processed during the study period, 466 (37.16%) samples yielded positive cultures and the rest had no growth.

466 culture positive cases yielded 512 isolates. Out of them, 306 (59.76%) were Gram negative bacteria, 196 (38.28%) were Gram positive bacteria and 10 (1.95%) were non albicans Candida.

Majority of the Gram positive bacteria were isolated from females i.e. 166 (84.69%). Majority i.e. 80 (40.81%) were from patients in the age group of 21-30 years followed by 24 (12.24%) in 31-40 years as shown in **Table 1**.

136 (69.38%) of Gram positive isolates were from outpatient department whereas 60 (30.61%) were from inpatient department as shown in **Table 1**.

Table 1: Different variables associated with Gram positive bacteria (n = 196)

Variables	Gram positive isolates	
	No.	%
Gender	Male	30 15.30
	Female	166 84.69
Age (years)	0-10	14 7.14
	11-20	22 12.79
	21-30	80 40.81
	31-40	24 12.24
	41-50	18 9.18
	51-60	22 11.22
	61-70	14 7.14
*OP/IP	OP	136 69.38
	IP	60 30.61

*OP- Out patient department

*IP- In patient department

Table 2 depicts the frequency of distribution of Gram positive bacteria from patients of various departments. Majority i.e. 132 (67.34%) were from the department of Obstetrics and Gynaecology followed by 28 (14.28%) from General Surgery, 16 (5.81%) from General Medicine, 14 (7.14%) from pediatrics and only 4 (2.32%) from orthopedics.

Table 2: Distribution of Gram positive bacteria according to various departments (n=196)

Department	No.	%
Medicine	16	5.81
OBG	132	67.34
Surgery	28	14.28
Orthopedics	4	2.32
Pediatrics	14	7.14


Table 3 shows the distribution of various Gram positive bacteria isolated from urine samples. Enterococcus 86 (43.87%) were the most common organisms isolated followed by Coagulase negative Staphylococcus (CONS) 80 (40.81%) and *Staphylococcus aureus* 30 (15.3%).

Table 3: Distribution of different Gram positive bacteria isolated from urine samples (n=196)

Organisms	No.	%
Enterococcus species	86	43.87
CONS (Coagulase negative Staphylococcus)	80	40.81
Staphylococcus aureus	30	15.3

Among the 196 Gram positive cocci, Methicillin sensitive *Staphylococcus aureus* (MSSA) were 18 (9.18%), Methicillin resistant *Staphylococcus aureus* (MRSA) were 12 (6.12%), Methicillin sensitive CONS were 36 (18.36%) and Methicillin resistant CONS were 44 (22.44%). *Enterococcus faecalis* accounted for 66 (33.67%) and other Enterococcus species 20 (10.2%) as depicted in **Fig. 1**

Fig. 1: Distribution of various Gram positive isolates from urine samples



Majority i.e. 126 (64.28%) of Gram positive cocci showed resistance to penicillin followed by resistance to norfloxacin in 100 (51.02%) as shown in **Table 4**.

Table 4: Antimicrobial susceptibility pattern of Gram positive isolates (n = 196).

Antibiotics	Susceptible		Resistant	
	No.	%	No.	%
Ampicillin	70	35.71	126	64.28
Tetracycline	134	68.36	62	31.63
Cotrimoxazole	42	21.42	72	36.73
Ciprofloxacin	120	61.22	76	36.2
Nitrofurantoin	184	93.87	12	6.12
Gentamicin*	146	74.48	50	26.12
Vancamycin	196	100	0	0
Linezolid	196	100	0	0
Norfloxacin	96	48.97	100	51.02

*High level gentamicin in case of Enterococcus species

All the Gram positive cocci were sensitive to Vancomycin and Linezolid. Majority of the Gram positive cocci were sensitive to nitrofurantoin with *Staphylococcus aureus* sensitivity of 28 (93.33%), CONS 78 (97.5%) and Enterococcus species 78 (90.69%). They were moderately sensitive to gentamicin (high level gentamicin in case of Enterococcus species) with *Staphylococcus aureus* sensitivity of 22 (73.33%), CONS 62 (77.5%) and Enterococcus species 62 (72.09%) as shown in **Table 5**.

Table 5: Antimicrobial susceptibility pattern of various Gram positive bacteria (n = 196).

Antibiotics	Staphylococcus aureus (n=30)		CONS (n=80)		Enterococcus species (n=86)	
	S	R	S	R	S	R
	No. %	No. %	No. %	No. %	No. %	No. %
Ampicillin	12 (40)	18 (60)	6 (7.5)	74 (92.5%)	52 (60.46)	34 (39.53)
Tetracycline	18 (45)	12 (40)	66 (82.5)	14 (17.5)	50 (58.13)	36 (41.86)
Cotrimoxazole	8 (26.66)	22 (73.33)	14 (17.5)	66 (82.5)	20 (23.25%)	66 (76.74%)
Ciprofloxacin	14 (46.66)	16 (53.33)	60 (75)	20 (25)	46 (53.48)	40 (46.51)
Nitrofurantoin	28 (93.33)	2 (6.66)	78 (97.5)	2 (2.5)	78 (90.69)	8 (9.30)
Gentamicin*	22 (73.33)	8 (26.66)	62 (77.5)	18 (22.5)	62 (72.09)	24 (27.9)
Norfloxacin	8 (26.66)	22 (73.33)	44 (55)	36 (45)	44 (51.16)	42 (48.83)
Vancomycin	30 (100)	0 (0)	80 (100)	0 (0)	86 (100)	0 (0)
Linezolid	30 (100)	0 (0)	80 (100)	0 (0)	86 (100)	0 (0)

*High level gentamicin in case of Enterococcus species

Discussion

The present study shows the pattern of UTIs prevalent in different age and sex groups, their causative organisms and the antimicrobial susceptibility pattern of Gram positive bacteria isolated. Out of the total 1254 urine samples processed, only 464 (37%) showed growth which is comparable with other studies done in different study settings and time periods.^{8,13-16} However, the proportion of positive cultures was higher in a couple of studies^{3,15} while few other studies reported low isolation rates.^{1,4,17,18} Further, 196 (38.28%) Gram positive bacteria were isolated in our study similar to the findings of other studies^{15,18} while lower rates were reported by other studies.^{4,14,17}

The proportion of females was more in patients presenting with symptoms of UTI i.e.166 (84.69%) which correlates well with the findings of other studies.^{3-5,17-20} Females are more prone to UTIs than men because of short urethra and its close proximity to the anus, which could be the reason for present study findings. On the contrary a study by Bajpai T et. al. reported higher prevalence of UTI in males.¹⁴ Majority i.e. 80 (40.81%) of the patients were in the age group of 21-40 years. Similar trend was reported by Kumar S. et. al.⁴

In our study, Enterococcus species was the commonest isolated Gram positive cocci 86 (43.87%) followed by CONS 80 (40.81%). This result is in agreement with previous studies.^{3,14,21-23} On the contrary some studies have isolated CONS^{4,15} and *Staphylococcus aureus*^{17,18} as the predominant uropathogens in their studies. This may be due to geographical differences in study settings, differences in practices related to hygiene and differences in

healthcare practices. Personal hygiene plays an important role in reducing the incidence of UTI as pathogens causing UTI are usual commensals in the perianal and vaginal regions.

All the Gram positive cocci were sensitive to Vancomycin and linezolid in the present study. This was in accordance with other studies.^{4,17,18,23,24} Majority of the Gram positive bacteria i.e. 93.88% were found to be sensitive to nitrofurantoin, similar to various studies done previously.^{17,20,23-26} and in contrast to some other studies which have observed a higher degree of resistance to nitrofurantoin.^{1,26} Owing to its narrow spectrum of activity, limited number of indications, limited tissue distribution and limited contact with bacteria outside the urinary tract, Nitrofurantoin has the highest activity among commonly used oral antibiotics for treatment of UTI.¹ On the other hand, the least effective drugs against Gram positive bacteria were ampicillin and norfloxacin with 64.28% and 51.02% of resistant strains respectively. Ampicillin resistance may be determined by the organisms due to the production of enzymes such as betalactamase.^{18,23,24,28-30}

Conclusion

Knowledge of the uropathogens and their antimicrobial susceptibility pattern in an area is essential for providing effective therapy and control of UTI. Antibiotic sensitivity pattern may vary even from place to place and from time to time, making periodical evaluation of antibiotic sensitivity a prerequisite for any hospital infection control program. The rise in the resistance pattern alerts us against indiscriminate usage of antibiotics and indicates the need to establish

antibiotic policies along with stringent measures to ensure effectiveness of the same.

References

- Mokta KK, Mokta JK, Verma S, Singh D, Kanga A. Bacterial etiology and antibiotic susceptibility pattern of urinary tract infection in sub-Himalayan region of India - a retrospective study of clinical isolates. *National Journal of Medical and Allied Sciences* 2015;4(1):38-45.
- Rajput MS, Chavan NS, Parashar S, Jain SM. Bacteriological profile of urinary tract infection in female patient attending tertiary care hospital, Indore, India. *Int.J.Curr.Microbiol.App.Sci* 2014;3(7):847-52.
- Moue A, AktaruzzamanSAQM, Ferdous N, Karim R, Khalil MMR, Das AK. Prevalence of urinary tract infection in both outpatient department and in patient department at a medical college setting of Bangladesh. *International Journal of Biosciences* 2015;7(5):146-152.
- Kumar S, Budhani D, Sayal P. Bacterial uropathogens and empirical treatment in urinary tract infection in a tertiary care institute. *Int.J.Curr.Microbiol.App.Sci* 2016;5(4):47-54.
- Angami S, Jamir N, Sarma PC, Deka AC. Urinary tract infection, its causative microorganism and antibiotic susceptibility in Nagaland. *Arch Med Health Sci* 2015;3(1):40-43.
- Hooton TM. Clinical practice. Uncomplicated urinary tract infection. *N Engl J Med*. 2012;366:1028-37.
- Wagenlehner FM, Naber KG. Current challenges in the treatment of complicated urinary tract infections and prostatitis. *Clin Microbiol Infect*. 12:67-80.
- Edwards MS, Baker CJ. Group B streptococcal infections in elderly adults. *Clin Infect Dis*. 2005;41:839-47.
- Mathews SJ, Lancaster JW. Urinary tract infections in the elderly population. *Am J Geriatr Pharmacother*. 2011;9:286-309.
- Baron EJ, Finegold SM. Microorganisms encountered in the urinary tract. In: Bailey & Scott's diagnostic microbiology. 9th edition. St. Louis, Missouri: Mosby publishers; 1994.
- Collee JG, Miles RS, Watt B. Test for the identification of bacteria. In: Collee JG, Fraser AG, Marmion BP, Simmons A (editors). Mackie and McCartney. Practical Medical Microbiology 14th ed. London: Churchill Livingstone 1996;131-145.
- Clinical and Laboratory Standard Institute. Performance standards for antimicrobial susceptibility testing; twenty-seventh Informational supplement. CLSI Document M100-S27 Wayne, PA. 2017;56-67.
- ManzoorKadri S, Gash B, Rukhsana A. Antibiotic sensitivity and resistance profile of the microorganisms responsible for urinary tract infection observed in Kashmir, India. *Ind medica* 2004;1(1):9-10.
- Bajpai T, Pandey M, Varma M, Bhatambare GS. Prevalence of extended spectrum beta-lactamase producing uropathogens and their antibiotic resistance profile in patients visiting a tertiary care hospital in central India: Implications on empiric therapy. *Indian J Pathol Microbiol* 2014;57(3):407-12.
- Singhal A, Sharma R, Jain M, Vyas L. Hospital and community isolates of uropathogens and their antibiotic sensitivity pattern from a tertiary care hospital in North West India. *Ann Med Health Sci Res* 2014;4(1):51-6.
- HariKattel P. Bacteriology of urinary tract infection among patients attending Tribuvan University teaching hospital Kathmandu, Nepal. *J. Nepal Assoc. Med. Lab. Sci*. 2008;9(1):25-9.
- Baral R, Timilsina S, Jha P, Bhattarai NR, Poudyal N, Gurung R, Khanal B, Bhattacharya SK. Study of antimicrobial susceptibility pattern of Gram positive organisms causing UTI in a tertiary care hospital in eastern region of Nepal. *Health Renaissance* 2013;11(2):119-24.
- Tayebi Z, Seyedjavadi SS, Goudarzi M, Rahimi MK, Boromandi S, Bostanabad SZ, Mirzaii A, Mahdiyoun M. Frequency and antibiotic resistance pattern in Gram positive uropathogens isolated from hospitalized patients with urinary tract infection in Tehran, Iran. *Journal of Genes, Microbes and Immunity* 2014;1-9.
- Katiyar R, Deorukhkar S, Siddiqui AU. Bacteriological profile and antibiogram of uropathogens with special reference to extended spectrum beta lactamases (ESBLs) detection in gram negative bacilli. *Indian Journal of Basic and Applied Medical Research* 2016;5(2):290-9.
- Shrestha P, Malla S, Basanta S, Dumre S, Upadhya B, Lamichhane S. Antimicrobial resistance pattern of bacterial isolates causing urinary tract infection. *JNHRC* 2002;5:49-54.
- Bajpai T, Pandey M, Varma M, Bhatambare GS. Mixed flora in the urine of hospitalized and elderly patients: Contamination or True infection?. *Niger J Exp Clin Biosci* 2014;2:20-7.
- Bhatnagar R, Soni P, Kumar S, Mehra SK, Solanki A and Tomar J. Study of nitrofurantoin susceptibility among uropathogenic bacterial isolates at a tertiary care hospital, Udaipur, Rajasthan. *World Journal of Pharmaceutical Research* 2015;4(5):2124-32.
- Preethishree P, Rai RK, Kumar V, Pai A and Bhat PU. Uropathogens and their antibiotic susceptibility pattern at a tertiary care teaching hospital in Coastal Karnataka, India. *Int.J.Curr.Microbiol.App.Sci*. 2016;5(1):23-31.
- Manjunath GN, Prakash R, Annam V, Shetty K. Changing trends in the spectrum of antimicrobial drug resistance pattern of uropathogens isolated from hospitals and community patients with urinary tract infections in Tumkur and Bangalore. *Int J Biol Med Res*. 2011;2(2):504-7.
- Jha VC, Yadav JN. Bacterial species isolated from urine of UTI suspected patients and their sensitivity to commonly available antibiotics. *J Nep Med Assoc* 1992;30:222-5.
- Levitt PN. Analysis of pathogens isolated from urinary tract infection in Barbados. *West Indi Med J* 1993;42:72-6.
- Grude N, Tveten Y, Krstiensen BE. Urinary tract infections in Norway: bacterial etiology and susceptibility. A retrospective study of clinical isolates. *Clin microbial infect* 2001;7(10):543-7.
- Forbes BA, Sahm DF, Weissfeld AS. Bailey and Scott's Diagnostic Microbiology 11th ed. USA: Mosby 2002.
- Raja MM, John SA. Multidrug resistance profile of urinary tract infected Gram positive pathogenic bacterial isolates. *Int J Infect*. 2015;2(1):22774.

How to cite this article: Naik TB, Lavanya J, Upadhya A, Mani V. Gram positive uropathogens and their antibiogram: Data analysis at a tertiary care hospital in Karnataka. *Indian J Microbiol Res* 2018;5(1):71-75.