

Neonatal septicemia: Blood culture bacterial isolates and their antimicrobial susceptibility pattern

Payal Soni¹, Mital Kumari Gamit^{2*}, Alka Nerukar³

^{1,2}Tutor, ³Professor and HOD, Dept. of Microbiology, ¹⁻³Gujarat Medical Education and Research Society Medical College and Hospital Valsad, Gujarat, India

*Corresponding Author: Mital Kumari Gamit

Email: gamit.mital@gmail.com

Abstract

Introduction: Neonatal septicemia is defined as “a clinical syndrome of bacteremia with systemic signs and symptoms of infection in the first four weeks of life”. Blood culture is the gold standard method for diagnosis of septicemia. Isolation of organism in blood culture and its antimicrobial sensitivity pattern carries a great role.

Aim: To provide prevalence and antimicrobial susceptibility pattern of bacteria isolated from suspected cases of neonatal septicemia at a tertiary care hospital, Valsad.

Materials and Methods: Retrospective study is conducted at a tertiary care hospital, Valsad from January 2017 to June 2018. Processing of blood culture samples, Isolation and identification of bacteria were done using standard microbiology techniques. Antimicrobial susceptibility testing was performed using modified Kirby-Bauer disk diffusion method as per Clinical laboratory Standard Institute guideline. ATCC E. coli 25922, ATCC S. aureus 25923, ATCC P. aeruginosa 27853 were used as standard strains.

Results: Out of 452 blood cultures, organisms were isolated in 127(28.09%). Commonly isolated bacteria are Coagulase Negative Staphylococci (42.52%) followed by Klebsiella spp. (18.11%), Acinetobacter spp. (15.75%), Enterococcus spp. (9.45%) and others (Staphylococcus aureus, Escherichia coli, Pseudomonas aeruginosa). Coagulase Negative Staphylococci strains have highest sensitivity to Vancomycin (100%), Linezolid (100%), Teicoplanin (100%), followed by Chloramphenicol (92.6%), Levofloxacin (90.7%), Cefoxitin (83.3%), and Tetracycline (77.8%). 16.7% isolates were Methicillin Resistant Staphylococci (MRS). Klebsiella spp. strain have highest sensitivity to Meropenem (73.9%), Imipenem (73.9%) and Amikacin (69.6%) followed by Chloramphenicol (65.2%), Tetracycline (60.9%) and Piperacillin Tazobactam (52.2%). 78.3% strains were Extended spectrum Beta-Lactamase (ESBL) producers.

Conclusion: Antimicrobial resistances are growing among bacteria and it differs according to different areas. So, proper surveillance system should be established to guide antimicrobial policy at local level.

Keywords: Neonatal septicemia, Blood culture, Antimicrobial susceptibility.

Introduction

Neonatal septicemia is defined as “a clinical syndrome of bacteremia with systemic signs and symptoms of infection in the first four weeks of life”.¹ Blood culture is the gold standard method for diagnosis of septicemia.²

According to WHO data, in 2016, Neonatal mortality rate (NMR) per 1000 live births globally is 18.6, whereas the same of South-East Asia region is 22.6 and of India is 25.4. Under five mortality rate among children of India is 43 per 1000 live births.³ So, developing countries like India still need proper measures to reduce it. Major causes of neonatal mortality worldwide are infections (36%, which includes sepsis/pneumonia, tetanus and diarrhoea), pre-term (28%), and birth asphyxia (23%).⁴ Neonatal septicemia is important factor responsible for morbidity as well as mortality. Isolation of organism in blood culture and its antimicrobial sensitivity pattern carries a great role.

Antimicrobial resistance among bacteria is growing world-wide and the same scenario exist even in developing countries like India.⁵ Proper surveillance system is required for better understanding of the situation and to provide guidance in making health policies at local, regional, national and international level, which is lacking in India.⁶

Aim of this study is to provide prevalence and antimicrobial sensitivity pattern of bacteria isolated from suspected cases of neonatal septicemia at tertiary care hospital, Valsad.

Materials and Methods

Retrospective study is conducted taking data of blood cultures received in microbiology laboratory at a tertiary care hospital, Valsad, Gujarat, India from suspected cases of neonatal septicemia from January 2017 to June 2018. Blood cultures were received in Brain-Heart Infusion broth, incubated aerobically at 37°C and repeated subcultures were done on Blood agar and MacConkey agar. Isolated organisms were identified using standard microbiology techniques.⁷ Antimicrobial sensitivity testing (AST) were performed on Muller Hinton agar using modified Kirby-Bauer disk diffusion method as per Clinical laboratory Standard Institute guideline 2017 and 2018.^{8,9} ATCC E. coli 25922, ATCC S. aureus 25923, ATCC P. aeruginosa 27853 were used as standard strains. Data was analyzed using Microsoft Excel 2016.

Results




Fig. 1: Result of culture

Table 1: Frequency and percentage positivity of bacterial isolates

Organisms	Frequency (out of 127)	Percentage positivity (%)
Coagulase Negative Staphylococci	54	42.52
Enterococcus spp.	12	9.45
Staphylococcus aureus	8	6.3
Bacillus subtilis	2	1.57
Klebsiella spp.	23	18.11
Acinetobacter spp.	20	15.75
E.coli	5	3.94
Pseudomonas aeruginosa	3	2.36

Table 2: Age wise distribution of samples

Age	Total	Positive	Percentage positivity
<8 days	367	99	26.97%
8 days or more	85	28	32.94%

(p value: 0.42 > 0.05, Difference is statistically not significant)

Table 3: Sex wise distribution of samples

Sex	Total	Positive	Percentage positivity
Female	185	51	27.56 %
Male	267	76	28.46 %

(p value: 0.87 > 0.05, Difference is statistically not significant)




Fig. 2: AST pattern of Coagulase Negative Staphylococci (n=54)




Fig. 3: AST pattern of klebsiella spp. (n=23)




Fig. 4: AST pattern of acinetobacter spp. (n=20)

Discussion

Result shows that the common blood culture bacterial isolates from NICU are Coagulase Negative Staphylococci and Klebsiella spp. with blood culture positivity rate of 28.09%.

Out of 452 blood cultures included in the study, organisms were isolated in 127(28.09%) as shown in Fig. 1. Frequency and percentage positivity of all isolated organisms is shown in Table: 1. Gram positive and Gram Negative bacterial isolates were 76(59.84%) and 51 (40.16%) respectively. Similar studies conducted by Hitesh J Assudani et al.¹⁰, Patel D et al.¹¹ and Shah Manisha et al.¹² in Gujarat shows positivity rate of 35.34%, 43.36% and 20% respectively with Klebsiella spp. and Coagulase Negative Staphylococci or Staphylococcus aureus as commonly isolated organism.

Positivity rate depend on many factors like non bacterial causes of septicemia, anaerobic bacterial infection, ongoing antibiotic therapy, less volume of blood etc.¹⁰ Klebsiella spp. and Coagulase Negative Staphylococci are commonly isolated organism in different studies^{14,18,19} may be because of their nature as saprophyte or commensals,²² environmental factors, low birth weight or prematurity which makes them more susceptible to surrounding infectons.^{10,23} Some studies also found E.coli and Staphylococcus aureus as commonly isolate organisms^{16,17,20,21} may be because of different geographical area.^{10,23}

Age and sex wise distribution is shown in Table: 2 and Table: 3. Difference is statistically not significant. (p value < 0.05).

Coagulase Negative Staphylococci strains have highest sensitivity to Vancomycin (100%), Linezolid (100%), Teicoplanin (100%), followed by Chloramphenicol (92.6%), Levofloxacin (90.7%), Cefoxitin (83.3%), and Tetracycline

(77.8%) as shown in Fig. 2. 16.7% isolates were Methicillin Resistant Staphylococci (MRS). Similar study conducted by Patel D et al.¹¹ shows 33% prevalence of MRS strains.

Klebsiella spp. strains have highest sensitivity to Meropenem (73.9%), Imipenem (73.9%) and Amikacin (69.6%) followed by Chloramphenicol (65.2%), Tetracycline (60.9%) and Piperacillin Tazobactam (52.2%) as shown in Fig. 3. All strains shows higher resistance to Cephalosporins. 78.3% strains were Extended Spectrum Beta-Lactamase (ESBL) producers. Similar study conducted by Patel D et al.¹¹, Hitesh J Assudani et al.¹⁰ and Shah Manisha et al.¹² shows 94.87%, 37.5% and 43.47% ESBL Klebsiella spp. prevalence respectively. Only 4 strains (17.39%) were Multi drug resistance. (MDR) Similar study conducted by Mamatha P. Samaga et al.¹³ shows 83.3 % MDR gram Negative bacteria.

Acinetobacter spp. strains shows highest sensitivity to Levofloxacin (100%) and Polymyxin B (100%) followed by Tetracycline (80%), Amikacin (70%), Piperacillin-Tazobactam (60%), Imipenem (60%) and Meropenem (60%). Different cephalosporins shows sensitivity varying from 20-40%.

All strains of Staphylococcus aureus (n=8) were sensitive to Cefoxitin, Teicoplanin, Linezolid and Vancomycin along with flouroquinolones. All strains of Enterococcus (n=12) were sensitive to Teicoplanin, Linezolid and Vancomycin.

Low prevalence of MRS strains and MDR organisms in our study suggest good hygienic practices among health care workers. Still lower susceptibility of Penicillin group and Cephalosporins among all bacteria create alarming sign for developing Anti Microbial resistance as they are commonly used drugs for Outpatient based treatment. Spread of these resistance strains leads to physical, financial and emotional burden to patients as well as community. There should be Antimicrobial usage guidelines based on local antimicrobial resistance data to reduce further development of resistance.

Conclusion

Klebsiella spp. and Coagulase Negative Staphylococci are common isolates in NICU. Resistance to frequently used Penicillin and Cephalosporin group is high among all bacterial isolates. Antimicrobial stewardship program should be implemented to make antibiotic policy at local level to deal the current scenario.

Conflicts of Interest: None.

References

1. Anand J. Amin et al., Sensitivity and resistance pattern of antimicrobial agents used in cases of neonatal sepsis at a tertiary care centre in western India, *Int J Pharm sci Res*, 2016;7(7):3060-3067.
2. Liyun Xu et al., Diagnostic value of C-Reactive Protein in neonatal sepsis: A meta analysis, *Eur j Inflamm* 2016;14(2) 100-108.
3. http://www.who.int/gho/child_health/mortality/neonatal/en/

4. http://www.who.int/pmnch/media/press_materials/fs/fs_newbornhealth_illness/en/
5. S. Ganesh kumar et al., Antimicrobial resistance in India: A review, *J Nat Sci Biol Med*, 2013;4(2):286-291.
6. BIRDY study group, www.thelancet.com, 387 February 6, 2016
7. Collee JG, Miles RS, Watt B. Tests for the identification of bacteria. In: Collee JG, Fraser AG, Marmion BP, Simmons A, editors. Mackie and McCartney Practical Medical Microbiology. 14th ed. Edinburgh: Churchill Livingstone; 1996.131-50.
8. M100, Performance Standards for Antimicrobial Susceptibility Testing, 27th edition, CLSI, 2017
9. M100, Performance Standards for Antimicrobial Susceptibility Testing, 28th edition, CLSI, 2018
10. Hitesh J Assudani et al., Bacteriological profiles of septicaemia in neonates at tertiary care hospital, Gujarat, *India J Res Med Dent Sci* 2015;3(2).
11. Patel D et al., Blood culture isolates in neonatal sepsis and their sensitivity in Anand District of India, *Int J Pharm Sci Res*, 2016;7(7):3060-3067.
12. Shah Manisha et al., Emergence of multi-drug-resistant Klebsiella pneumoniae in Neonatal Intensive Care Units: concern about antimicrobial policies, *Res J Recent Sci* 2012;1(ISC-2011),275-280.
13. Mamatha P. Samaga et al., Prevalence of neonatal septicaemia in a tertiary care hospital in Mandya, Karnataka, India, *Int J Res Med Sci* July 2016;4(7):2812-2816.
14. Shilpa Pradhan et al., Blood Culture of Neonates in Paediatric Department and their Antimicrobial Susceptibility Pattern in and around Nims University, Jaipur, India, *Int J Curr Microbiol Appl Sci* 2017;6(4):1940-1946.
15. Rajlakshmi Viswanathan et al., Profile of Neonatal Septicaemia at a District-level Sick Newborn Care Unit, *J Health Popul Nutr* 2012;30(1):41-48.
16. Dechen Tsering et al., Bacteriological profile of septicemia and the risk factors in neonates and infants in Sikkim, *J glob infect dis* 2011;3(1):42-45.
17. Vrishali Avinash Muley et al., Bacteriological Profile of Neonatal Septicemia in a Tertiary Care Hospital from Western India, *J glob infect dis* :2015;7(2):75-77.
18. P. Jyothi et al., Bacteriological profile of neonatal septicemia and antibiotic susceptibility pattern of the isolates, *J Natural Sci Biol Med* 2013;4(2): 306-309.
19. Sneha Ann Oommen et al., Bacteriological Profile Of Neonatal Septicemia: A Retrospective Analysis From a Tertiary Care Hospital In Loni, *Int J Med Res Health Sci* 2015;4(3):652-658.
20. Subedi D et al., Bacteriological profile of Neonatal septicemia, <https://www.researchgate.net/publication/265382116>
21. Shrestha NJ et al., Bacteriological Profile of Neonatal Sepsis: A Hospital Based Study, *J Nepal Paediatr Soc* 2011;31(1).
22. Essentials of Medical Microbiology, Apurba Shankar Sastry, Edition 1, page no: 219, 308
23. Shaw CK, Shaw P, Thapalia A. Neonatal sepsis bacterial isolates and antibiotic susceptibility pattern at a NICU in a tertiary care hospital in western Nepal: A retrospective analysis. *Kathmandu Univ Med J* 2007;5(2):153-60.

How to cite this article: Soni P, Gamit MK, Nerukar A, Neonatal septicemia: Blood culture bacterial isolates and their antimicrobial susceptibility pattern, *Int J Med Microbiol Trop Dis* 2019;5(1):14-17