



Review Article

Quorum sensing — Polymicrobial challenge in periodontics

Sangeetha S¹, Padmalochani B², Jaishree Tukaram Kshirsagar¹,
Sai Sarath Kumar Kothimbakkam³, Burnice Nalina Kumari C^{4,*}

¹Dept. of Periodontics, Tamilnadu Government dental college and hospital, Tamil Nadu, Chennai, India

²Dept. of Periodontist, Abeer Alnoor Polyclinic, Muhaisnah, Dubai, United Arab Emirates

³Ragas Dental College and Hospital, Tamil Nadu, Chennai, India

⁴Dept. of Periodontics, Meenakshi Ammal Dental College and Hospital, Tamil Nadu, Chennai, India



ARTICLE INFO

Article history:

Received 01-12-2019

Accepted 22-12-2019

Available online 07-01-2020

Keywords:

Quorum sensing

Quorum quenching

Bacterial communication

ABSTRACT

Periodontal disease is a multifactorial disease caused by imbalance between complex microbial community and the host immune response. Plaque provides favorable environment for the survival and growth of microorganisms inducing periodontal disease. The current concept of biofilm formation and cell to cell communication of microorganisms present in the biofilm occurs by the production of chemical signaling molecules which has become the niche for molecular level research projects. This biological process is called quorum sensing which modulates the expression of genes involved in survival, biofilm formation, virulence and pathogenicity of microorganisms. To eradicate this communication, quorum quenchers play a role in the inhibition of signals of quorum sensing. This review article enlightens the quorum sensing signaling and the magical action of quorum quenchers from the data collected by different analysis.

© 2019 Published by Innovative Publication. This is an open access article under the CC BY-NC-ND license (<https://creativecommons.org/licenses/by/4.0/>)

1. Introduction

Periodontal disease is an inflammatory disease which is multifactorial in origin causing destruction of the periodontium and also acts as a risk factor for systemic diseases such as cardiovascular diseases, endocrine disorders, hemorrhagic disorders, renal diseases, liver diseases, pulmonary diseases, and pre-term low birth weight infants.¹ Throughout life, body interface surfaces are exposed to microbial colonization resulting in establishment of microbiota lives in harmony with the host. As teeth provide hard, non-shedding surfaces there is a development of extensive bacterial deposits. The accumulation and metabolism induced growth of bacteria on tooth surfaces is the influencing factor for initiation of gingivitis and periodontitis.

On mechanical cleaning of teeth, hydrophobic macromolecules begin to adsorb to the tooth surface forming

conditioning film known as acquired pellicle which consists of various salivary glycoproteins and antibodies. This results in alteration of the surface free energy which increases the efficiency of bacterial adhesion. Bacteria adhere to these coated surfaces leading to formation of a biofilm. In 1mm² of dental plaque weighing approximately 1 mg contains 200 million bacteria.

1.1. Plaque Formation

Approximately 700 species of microorganisms colonize the human oral cavity which includes commensals and sparse population of pathogenic bacteria. Primary colonization of bacteria occurs by adsorbing on to the pellicle coated surface. Facultative anaerobic gram – positive cocci dominate. At 24 hours plaque mainly shows streptococci, of which *Streptococcus sanguis* is predominant. Streptococci and other precursor organisms provide unique receptor sites for later, more pathogenic colonizers such as *Fusobacterium nucleatum* which acts

* Corresponding author.

E-mail address: dr_burnice@yahoo.co.in (B. N. Kumari C).

as bridging bacteria, *Tannerella forsythia*, *Treponema denticola*, and *Porphyromonas gingivalis*.³ which are closely associated with the development of periodontitis.²

F. nucleatum, *P. intermedia*, and *P. gingivalis* are the three very important microorganisms in the development of periodontal disease. *Fusobacterium nucleatum* plays an important role in the subgingival microbiota. It has ability to coaggregate with most of the microorganisms in plaque and with periodontal pathogens. *P. intermedia* is elevated in acute necrotizing ulcerative gingivitis. *P. gingivalis* plays an important role in bone and tissue destruction.

Micro-organisms are mostly organized in communities and they assemble on substrates known as biofilm. A typical biofilm forms at the interface and comprises of microbial cells enclosed within a matrix consisting of polysaccharides, proteins, nucleic acid and lipids, derived from microbial and environmental sources.⁴ Plaque biofilm is the etiologic factor for periodontitis. This biofilm is stabilized by extracellular DNA (eDNA).⁵ In addition, this eDNA may be a source for potential transfer of antibiotic resistance or virulence genes between species within the communities.⁶ The creation of bio-films is regulated by signaling process called quorum sensing.

1.2. Quorum Sensing

Quorum is a Latin word. Bacteria produce chemical signals and other bacteria can respond to them by a mechanism through which they communicate population behavior and phenotype known as cell-cell communication or cell-cell Signaling or quorum sensing.^{7,8} As communication is the resolving factor to interpret discrepancies, biofilms forming bacteria adopt specialized roles and communicate with one another.⁹

To initialize communication between them, Autoinducers or pheromones which are small hormone like chemical signaling molecules play an important role. In formation and maturation of plaque. It either diffuses freely across the cell membranes or is actively transported out of the cell.¹⁰ Autoinducer 2 is described as a global signal molecule for interspecies communication, as it is made by gram-positive as well as gram-negative bacteria.¹⁰ When the population of Quorum Sensing producing bacteria leaps, the individual organisms engenders and secrete the autoinducers into the extracellular environment. Thus, the concentration of external autoinducer is congruent with cell population density. By monitoring the extracellular autoinducer concentration, the bacteria can count one another and alter target gene expression.¹¹ Though planktonic cells secrete chemical signals at low concentration, there is no change in genetic expression. When Biofilm cells held together in dense populations, the secreted chemical signals attain at higher concentrations. These molecules then re-cross the cell membranes and trigger changes in genetic activity leading to the production of various products which

increase bacterial pathogenicity⁹ (Table 1).

At this instance, bacteria produce a light inducing property to attract one another to increase dense population known as bioluminescence. The bacterial signaling was first described in the bioluminescent marine organism *Vibrio fischeri* [Nealson and Hastings, 1979] where a diffusible signal N-acyl homoserine lactone (AHL) was responsible for the induction of bioluminescence.¹⁶ In the light organ of the Hawaiian squid *Euprymna scolopes*, *V. fischeri* colonizes and induce the expression of genes required for bioluminescence. This bioluminescence increases the density of microorganism present in a particular area and makes cell to cell communication effective. When they are at high cell concentrations, the level of the autoinducer becomes adequate to induce transcription of the genes that produce the enzyme luciferase.¹⁶ This enzyme causes the oxidation of the reduced flavin mononucleotide to produce a long-chain fatty acid, water and flavin mononucleotide. This reaction causes the emission of blue-green light along with the oxidation reaction and therefore is termed as bioluminescence. Different luminescent bacteria may show different luminescence spectrum and colour of the emitted light due to the shift in wavelength caused because of the sensitizer proteins.¹⁷

Quorum Sensing is believed to regulate and control the traits of micro-organism such as competence development, sporulation, Antibiotic resistance, virulence factor, induction, Cell differentiation and nutrient flux along with other physiological events in pathogenic bacterial infections.^{11,18}

Finally, when the threshold concentration is reached, the population is considered to be "quorate."¹⁸ This causes the binding of the autoinducer to the cognate receptors present within the bacteria. This then further triggers a signal transduction cascade that results in wide-spread changes in the gene expression.¹⁹

1.3. Types of quorum sensing systems

Gram-negative bacteria: uses 2 proteins which controls the expression of luciferase operon (luxICDABE) required for light production.

1. The LuxI -type (auto inducer synthase) -involves in synthesis and emission of acyl homoserine lactone
2. The LuxR type (cytoplasmic autoinducer receptor)- binds to received acyl homoserine lactone.^{20,21}

After production, the AHL freely diffuses inner and outer membrane of the cell and increases its concentration in response to increasing cell density.²² When the signal reaches a critical, threshold concentration, it is bound by LuxR and this complex activates transcription of the operon encoding luciferase.²³ LuxR - AHL complex also acts as positive feedback activator, leading to high expression of luxI gene and production of LuxI protein.

Table 1: Autoinducers

Acylated homoserine lactone	Gram negative and Intraspecies
Peptides	Gram positive and Intraspecies
Furanosyl borate diester	Gram positive and gram negative, Interspecies

Table 2: Quorum Quenching compounds in plants

Garlic	Ajoene, Disulphides, Trisulphides ¹²
Turmeric	Curcumin ¹²
Citrus flavonoids	Flavonine naringenin ¹²
Horseradish	Iberin ¹²
Red Marine Algae(Dalea Pulchra)	Halogenated Furanones ¹³
Grape Fruit Extract	Furocoumarins, Carotenoids, Limonoids, Pectin ¹³
Clove Extract	Eugenol, Hexane, Menthol ¹⁴
Coffee Extract	Caffeine ¹⁵

BIOFILM DEVELOPMENT




Fig. 1: Biofilm formation




Fig. 2: Action of autoinducers

This regulation induces the environment with the signal production and produce light.²⁴ A large number of other gram-negative proteobacteria possess LuxIR-type proteins and communicate with AHL signals.²⁵ These systems are used predominantly for intra-species communication as extreme specificity exists between the LuxR proteins and their cognate AHL signals.²¹ The following figure shows the cell density dependent action of AHL.

Gram-positive bacteria use oligopeptides processed from precursors as autoinducers. These are auto inducing peptides:²⁶ Gram-positive bacteria use two processes in quorum sensing.²⁰ They are:

1. A two component signal transduction system
2. Internalization

1.4. Why there is a need for eradicating the quorum sensing process

Biofilms are more resilient to mechanical removal and to killing by the host immune system because the matrix within the biofilm protects bacteria from exposure to innate immune defenses and antibiotic treatments.²⁷ Hence, a new era of therapeutics have evolved which turned into a promising potential tool for abolition of infectious diseases caused by bacteria. Thus, Quorum sensing plays a key role in pathogenesis of disease causing by bacteria with its influence at the genetic level leading to regulation of various factors involved in the process.

1.5. Mode of Eradicating

Several ways are there for disrupting the quorum sensing as quorum sensing forms the basis of infection. Following ways are enlisted below:

1. Early colonizers are responsible for creating the pathogenic state. The Presence of Periodontal Pathogens in dental plaque could modulate the virulence properties of *S.mutans* By Interfering with Its Com Quorum Sensing System. Since The Com Quorum Sensing System Exists In Many Species of Earlier Dental Plaque Colonizers. This Interference of Quorum Sensing By Periodontal Pathogens Such As *P.gingivalis* and *T. denticola* Could, At Least In Part, Be A Mechanism of Bacterial Antagonism In Periodontal Diseases.¹
2. Quorum quenchers – synthetic and naturally occurring compounds.

1.6. Quorum quenchers-action of magic bullets

Prevention and control of bacterial infection by quorum quenching, which is an antipathogenic or signal interference of quorum sensing has been explored.²⁸ It retards the

action of quorum sensing by enzymatic hydrolysis of AHL autoinducers. Quorum sensing blockage are at different site.

1. QQ by small quorum-sensing inhibitors (QSI).
2. QQ by AHL- lactonase.
3. QQ by AHL- acylase.
4. QQ by paraoxonase enzymes²⁴

1.7. Natural quorum sensing inhibitors

1.7.1. Algae

Anti QS agents were first characterized in the marine red alga, *Delsia puchra*. Halogenated furanones produced by *Delsia puchra* could inhibit QS in a number of bacteria.

1.7.2. Fungi

Patulin and penicillic acid produced by *Penicillium coprobium* and *P. radicola*, respectively, as inhibitors of quorum sensing in *P. aeruginosa*. These two compounds target the Las R and RhIR QS regulators²⁹

1.7.3. Insects

A venom alkaloid from the fire ant *Solenopsis invicta*, solenopsin A - inhibits quorum sensing in *P. aeruginosa*³⁰

Honey had QSI activity against *Erwinia carotovora*, *Yersinia enterocolitica*, *Aeromonas hydrophilia*³¹ (Table 2)

1.8. Synthetic quorum sensing inhibitors

There are basically three ways to block QS by developing on the AHL scaffold²⁴

1. Introduction of substitutions in the acyl side chain without any change in the lactone ring.
2. Introduction of substitutions and alterations in the lactone ring with unchanged acyl side chain.
3. Extensive modifications in both the acyl side chain and the lactone ring.

The synthetic drugs which act as quorum sensing inhibitors are Macrolides, RNA III- related compounds, Halogenated furanones, Homoserine lactone analogue - Antagonism of Homoserine lactone activity and can be prepared as homoserine lactone vaccine.³²

2. Discussion

The biofilm formation can be disrupted by the naturally occurring quorum sensing inhibitors which plays pivotal as quorum quenchers in inhibiting formation of dental plaque and further progression of periodontal disease. The discovery and further research of cell signaling among microbes results as a new milestone in eradication polymicrobial disease.

3. Conclusion

Several studies are being done on the benefits of a quorum sensing inhibition. Understanding the extent and significance of bacterial intercellular communication is still in its early stages and more discoveries are required to know the real extent and significance of bacterial cell–cell communication in the environment. Phytochemical quorum sensing inhibitor compounds may serve as a panacea in dentistry. Further research should be carried out to study the ability of quorum quenchers.

4. Source of funding

None.

Acknowledgment and conflict of interest

None.

References

- Wang BY, Alvarez P, Hong J. HK Kuramitsu Periodontal pathogens interfere with quorum-sensing-dependent virulence properties in *Streptococcus mutans*. *J Periodontol Res*. 2011;46(1):105–110.
- Haffajee AD, Socransky SS. Microbial etiological agents of destructive periodontal diseases. *Periodontol*. 1994;5:78–111.
- Kolenbrander PE, Palmer RJ, Periswamy S, Jakubovics N. Oral multispecies biofilm development and the key role of cell-cell distance. *Nat Rev Microbiol*. 2010;8(7):471–480.
- Flemming HC, Wingender J. The biofilm matrix. *Nat Rev*. 2010;9(9):623–633.
- Aaron BMT, Katie BS, Rachel LS, Carol WL, Gary DM. Enterococcus faecalis produces abundant extracellular structures containing DNA in the absence of cell lysis during early biofilm formation. 2012; Available from: [10.1128/mBio.00193-1224](https://doi.org/10.1128/mBio.00193-1224).
- Roberts AP, Mullan P. Oral Biofilms: a reservoir of transferable, bacterial, antimicrobial resistance. *Expert Rev Anti Infect Ther*. 2010;8(12):1441–1450.
- Bhardwaj AK, Vinothkumar K, Rajpara N. Bacterial quorum sensing inhibitors: attractive alternatives for control of infectious pathogens showing multiple drug resistance. *Recent Pat Antiinfect Drug Discov*. 2013;8(1):68–83.
- Bassler Interspecies communication in bacteria. *Invest J Clin*. 2003;112:1291–1299.
- Mahajan A, Singh B, Kashyap D, Kumar A, Mahajan P. Interspecies Communication and Periodontal Disease. *Scientific World Journal*. 2013;p. 765434–765434.
- Baswaraj B, Prapulla. Quorum Sensing in Plaque Biofilms: Challenges and Future Prospects. *Dev*. 2011;p. 10024–1080.
- Surette MG, Miller MB, Bassler BL. Quorum sensing in *Escherichia coli*, *Salmonella typhimurium*, and *Vibrio harveyi*: A new family of genes responsible for autoinducer production. *Proc Natl Acad Sci*. 1999;96(4):1639–1644.
- Esterela AB, Abraham WR. Combining biofilm-controlling compounds and antibiotics as a promising new way to control biofilm infections. *Pharmaceuticals*. 2010;3:1374–1393.
- Koh CL, Sam CK, Yin WF. Plant-derived natural products as sources of anti-quorum sensing compounds. *Sensors*. 2013;p. 136217–136228.
- Zhou L, Zheng H, Tang Y, Gong Q. Eugenol inhibits quorum sensing at sub-inhibitory concentrations. *Biotechnol Lett*. 2013;3:631–637.
- Aparna Y, Sarada J. Quorum Quenchers - Past, Present and Future of This Novel Therapeutics. *Wjpls*. 2015;1(3):108–128.
- Nealson KH, Hastings JW. Bacterial Bioluminescence: Its Control and Ecological Significance. *Microbiological Reviews*. 1979;43(4):496–518.
- Bassler BL, Curr O. How bacteria talk to each other: Regulation of gene expression by quorum sensing. *Microbiol*. 1999;2:582–587.
- Suchetha A, Chitra J, Darshan BM, Sapna N, Apoorva SM, Nanditha C. Quorum sensing - Meetings in the Microbial World. *J Res Med Den Sci*. 2015;3(3):161–165.
- Meighen E. Encyclopedia of Microbiology. Academic Press ; 1992., p. 309–311.
- Kamalesh B, Sonwane S, Guptha I, Swetha RK. Quorum Sensing Inhibition, Relevance to Periodontics. *J Int Oral Health*. 2015;7(1):67–69.
- Christopher WM, Bonnie BL. Annu Quorum Sensing: Cell-to-Cell Communication in Bacteria. *Rev Cell Dev Biol*. 2005;21:319–346.
- Kaplan BH, P GE. Diffusion of Autoinducer is involved in regulation of the *Vibrio fischeri* Luminescence System. *Journal of Bacteriology*. 1985;p. 1210–1214.
- Stevens AM, Dolan KM, Greenberg EP. Synergistic binding of the *Vibrio fischeri* LuxR transcriptional activator domain and RNA polymerase to the lux promoter region. *Proc Natl Acad Sci*. 1994;91:12619–12623.
- Nealson KH, Hastings JW. Bacterial bioluminescence: Its control and ecological significance. *Microbiol Rev*. 1979;43:496–518.
- Turner SL. Quorum sensing in context: out of molecular biology and into microbial ecology. *Microbiology*. 2002;p. 3762–3764. Pt. 12.
- Sifri C. Healthcare epidemiology: quorum sensing: bacteria talk sense. *Clin Infect Dis*. 2008;47(8):1070–1076. Sifri CD.
- Leid JG, Willson CJ, Shirtliff ME, Hassett DJ, Parsek MR, Jeffers AK. The exopolysaccharide alginate protects *Pseudomonas aeruginosa* biofilm bacteria from IFN- γ -mediated macrophage killing. *J Immunol*. 2005;75(11):7512–7520.
- Dong YH, Wang LH, Xu JL, Zhang HB, Zhang XF, Zhang LH. Quenching quorum-sensing-dependent bacterial infection by an N-acyl homoserine lactonase. *Nature*. 2001;411(6839):813–817.
- Rasmussen TB, Bijarnsholt T, Skindersoe ME, Hentzer M, Kristoffersen P, et al. Screening for quorum-sensing inhibitors (QSI) by use of a novel genetic system, the QSI selector. *J Bacteriol*. 2005;187(5):1799–1814.
- Kim C, Kim J, Park HY, Park HJ, Lee JH, et al. Furanone derivatives as quorum-sensing antagonists of *Pseudomonas aeruginosa*. *Appl Microbial Biotechnol*. 2008;80(1):37–47.
- Truchado P, Gil A, Tomas FA, Barberan, Allende A. Inhibition by chestnut honey of N-acyl-l-homoserine lactones and biofilm formation in *Erwinia carotovora*, *Yersinia enterocolitica* and *Aeromonas hydrophila*. *Journal Agricultural and Food Chemistry*. 2009;57(23):11186–1193.
- Rasmussen TB, Givskov M. Quorum sensing inhibitors: a bargain of effects. *Microbiology*. 2006;152:895–904.

Author biography

Sangeetha S 2ND Year MDS

Padmalochani B Private Practitioner

Jaishree Tukaram Kshirsagar Professor

Sai Sarath Kumar Kothimbakkam 3rd Year MDS

Burnice Nalina Kumari C Assistant Professor

Cite this article: Sangeetha S , Padmalochani B , Kshirsagar JT, Kothimbakkam SSK, Kumari C BN. Quorum sensing — Polymicrobial challenge in periodontics. *Int J Periodontol Implantol* 2019;4(4):115-119.