

Periodontitis and liver disease a relation - Review

Shaik Ali Hassan

Dental surgeon, Dept. of Oral Medicine, Manav Rachna Dental College, Faridabad, Haryana, India

***Corresponding Author: Shaik Ali Hassan**

Email: alishaikhassan@gmail.com

Abstract

Periodontitis is an oral infection that is exceptionally common around the world, with a commonness of 30-50% of the populace in created nations, yet just 10% present with serious structures. It is additionally assessed that periodontitis brings about overall profitability misfortunes adding up to 54 billion USD yearly. Expanding proof additionally demonstrated that periodontitis may take an interest in the movement of liver illnesses, for example, non-alcoholic greasy liver sickness, cirrhosis and hepatocellular carcinoma, just as influencing liver transplantation. The aim of the review is that periodontitis may be important in the progression of liver disease, thus providing dentists and physicians with an improved understanding of this issue.

Keywords: Periodontitis, Liver disease, NAFLD, ALT, DM and CVDs.

Introduction

Periodontal diseases, which incorporate gum disease (where the inflammation is restricted to the gingiva and is reversible with dental consideration) and periodontitis (where the inflammation spreads, furthermore, brings about tissue obliteration and alveolar bone resorption), are the most widely recognized kinds of disease in people around the world.¹ Tissue devastation, for example, the breakdown of the collagen filaments of the periodontal tendon (PDL) and the loss of gingival tissue furthermore, alveolar bone, are normal for periodontitis, coming about in continuous debilitating of the tooth-supporting tissues, which in the end prompts tooth misfortune. The condition is broad. Extreme periodontitis that undermines tooth maintenance influences 10-15% of grown-ups in most of populaces examined also, went from 1%, among 20-to 29-year-old, to 39%, among people >65-year-old while moderate periodontitis influences 40-60% of grown-ups in all populaces.²

Periodontitis and liver disease

Roughly one-fourth of the whole grown-up populace on the world appears unreasonable hepatic fat gathering, and NAFLD (non alcoholic fatty liver disease) is the most basic type of incessant liver infection experienced in created nations.³ The

predominance of NAFLD all through the world is theorized to be 20-30%,⁴ among hefty patients the figure ascends to 57-74%.⁵ The conclusion of NAFLD is frequently settled after recognizable proof of raised serum alanine aminotransferase (ALT) and γ -glutamyl transferase (GGT), which is most generally utilized for screening of liver sicknesses in corpulent and asymptomatic patients.⁶

Interaction between NAFLD and periodontitis

In an examination directed in a Japanese college, male understudies with an elevated level of serum ALT were distinguished to be fundamentally bound to have periodontitis than those with a low degree of serum ALT.⁷ With respect to females, the relationship between a higher ALT level what's more, an expanded danger of periodontitis was not seen as huge, which was as opposed to a past report that showed that the occurrence pace of periodontitis in females matured 20-59 years was fundamentally expanded with raised serum levels of ALT.⁸ The creator of the previously mentioned examinations ascribed this disparity to the distinctions in the age of the subjects and the example size. Outstandingly, it appears that the relationship among periodontitis and the serum levels of ALT is shared. In a cross-sectional examination with an enormous example size that was directed in Japan,

scientists found that ALT and GGT levels were higher in patients with periodontal pockets (depth, ≥ 4 mm) when contrasted with solid controls. Various calculated relapse investigation with GGT or ALT as the ward variable uncovered that there was a critical affiliation between periodontal pockets and GGT, significantly in the wake of changing for age, sexual orientation, cigarette smoking, liquor drinking propensities, furthermore, side effects of MS (metabolic syndrome).⁹

Mechanism of periodontitis on liver disease

The microbial etiology of periodontal disease has been the focal point of research for quite a while. Around 400 species have been recognized in the gingival sulcus, among them are *Porphyromonas gingivalis* (*P. gingivalis*) and *Tannerella forsythia*, which are generally viewed as significant pathogens in periodontitis.¹⁰ Subgingival microbiota were characterized into a few buildings demonstrated by different hues; the hues (changing from red to yellow) have extraordinary implications, with red being the most pathogenic and yellow being less intrusive. Periodontal microbiota are more heterogeneous than prior accepted. In dentistry, gram-negative life forms were viewed as the dominating microorganisms in periodontitis; be that as it may, gram-positive life forms found in profound, diseased locales are proposed to be the most significant pathogens in periodontitis.¹¹ Microbes additionally have a negative impact on the liver. It is outstanding that patients with cirrhosis are at more serious danger of bacterial disease¹² and contaminations rate is 4-to 5-fold higher than the overall public.¹³

P. gingivalis is discharged from the sulcus into the circulatory system. Human preliminaries and creature tests have affirmed the nearness of *P. gingivalis* in liver tissues.¹⁴ Besides, the periapical granuloma, which filled in as a steady and feasible stockpile wellspring of the *P. gingivalis* and its items, may prompt incessant liver damage.¹⁵ In an investigation where the occurrence pace of *P. gingivalis* was thought about between NAFLD patients and non-NAFLD control subjects, it was discovered that the discovery recurrence of the *P. gingivalis* disease in NAFLD patients was fundamentally higher.⁸ *P. gingivalis*

disease perhaps a free indicator for the improvement of NAFLD and may add to the movement of other LD.

Aggregatibacter actinomycetemcomitans (*A. actinomycetemcomitans*). *A. actinomycetemcomitans* is an exogenous bacteria, which is related with periodontitis in young people and can create destructiveness factors.¹⁶ Studies have indicated that *A. actinomycetemcomitans* creates certain items, which may inactivate and dodge resistant safeguard. The most examined results of *A. actinomycetemcomitans* are leukotoxin and rehashes in poison.¹⁷ A past study demonstrated that the infusion of *A. actinomycetemcomitans* into mice prompted immunosuppression and smothered the IgG reaction to red platelets.¹⁸ The organization of *A. actinomycetemcomitans* has likewise been accounted for to prompt foundational irritation in apolipoprotein E-deficient mice.¹⁹ In a creature study, *A. actinomycetemcomitans* was available in liver tissue after intravenously immunizing mice with live *A. actinomycetemcomitans*. also, may also initiate moderate hepatic inflammation.

Inflammatory mediators

Inflammatory mediators, for example, IL-12/23, TNF- α , and IL-1 may prompt the enrollment of actuated neutrophils, which causes hepatocyte and vascular endothelial cell wounds by discharging oxidants and proteases.²⁰ Human preliminaries²¹ and animal tests²² have affirmed the generation of pro-inflammatory atoms in vivo and ex vivo in patients and creatures with cirrhosis. The liver goes about as the body's first safeguard against microorganisms and microbial segments. These pathogens, for example, inflammatory mediators, which are available in the entry blood, produce the underlying immunological and hormonal weight to the liver.²³

Cytokines and chemokines- The dental plaque, mostly made out of gram-negative microbes cell dividers, which are shaped of peptidoglycans, polysaccharides, proteins, lipids, lipopolysaccharides (LPSs) and lipoproteins,²⁴ ordinarily exist in the oral depression of people, especially in the individuals who experience the ill effects of periodontitis. Invigorated by these segments, the periodontal tissue produces

inflammatory cytokines, (for example, IL-1 β , IL-12, IL-10, IL-6, TNF- α and INF- γ) and chemokines [such as monocyte chemotactic protein 5 (MCP-5), IL-8 and macrophage inflammatory protein-1 α (MIP-1 α), prostaglandin E2 and nitric oxide (NO)].²⁵ These pro-inflammatory cytokines are engaged with the movement of LD, for example, cirrhosis.²⁶ Oral microscopic organisms are likewise significant in the cytokine arrange. LPSs, discharged by periodontal microscopic organisms, for example, *A. actinomycetemcomitans* and *P. gingivalis*, influence the safe framework by official to Toll-like receptor (TLR)-4 or -2, oral microbes additionally invigorate the declaration of co-stimulatory particles, bunch of separation (CD) 80/CD86 by authoritative to TLR4; and may take an interest in the enactment of T-cells and fuel liver irritation.²⁷ Kupffer cells, which express the most elevated levels of TLR4 in the liver, are the essential cells in liver aggravation that react to LPSs so as to deliver inflammatory cytokines, chemokines and responsive oxygen species (ROS).²⁸

The peptidoglycans- segments of the bacterial dividers,²⁹ of flowing oral microscopic organisms may invigorate platelets to deliver cytokines. Likewise with LPS, peptidoglycans add to the actuation of resistant cells by official to the TLR2 receptor.³⁰ Moreover, peptidoglycans can be perceived by the supplement framework and explicit receptors, consequently bringing about the creation of TNF- α , IL-6, IL-8, IL-1 β , MIP-1 α and NO in macrophages.³⁰ It has likewise been indicated that the levels of IL-6 explicitly expanded altogether following scaling (a dental practice frequently used to kill analytics), while the IL-8 levels diminished.³¹

Conclusion

Relationship among periodontitis and NAFLD, LC, HCC, and LT have recently been examined. Certain obsessive highlights are shared by periodontitis and foundational illnesses, for example, DM and CVDs, may apply comparative consequences for the liver. Among the three instruments (counting microscopic organisms, pro-inflammatory mediators and oxidative pressure), different microorganisms exist in the dental

plaque, some of which are progressively prevailing in patients with extreme periodontitis, may essentially add to the connecting of other neurotic components. So by this we conclude that there is relationship between the liver disease and periodontitis.

Source of Funding

None.

Conflict of Interest

None.

Reference

1. Vos T, Flaxman AD, Naghavi M, Lozano R, Michaud C, Ezzati M et al, Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012;380:2163-96.
2. Chung JH, Hwang HJ, Kim SH and Kim TH: Associations Between Periodontitis and Chronic Obstructive Pulmonary Disease; the 2010-2012 Korean National Health and Nutrition Examination Survey (KNHANES). *J Periodontol* 2016;25.
3. Angulo P: Nonalcoholic fatty liver disease. *N Engl J Med* 2002;346:1221-31.
4. Angulo P and Lindor KD: Non-alcoholic fatty liver disease. *J Gastroenterol Hepatol* 2002;17:S186-S190.
5. Milić S and Stimac D: Nonalcoholic fatty liver disease/steatohepatitis: Epidemiology, pathogenesis, clinical presentation and treatment. *Dig Dis* 2012;30:158-62.
6. Oh HJ, Kim TH, Sohn YW, Kim YS, Oh YR, Cho EY et al, Association of serum alanine aminotransferase and γ -glutamyltransferase levels within the reference range with metabolic syndrome and nonalcoholic fatty liver disease. *Korean J Hepatol* 2011;17:27-36.
7. Furuta M, Ekuni D, Yamamoto T, Irie K, Koyama R, Sanbe T et al, Relationship between periodontitis and hepatic abnormalities in young adults. *Acta Odontol Scand* 2010;68:27-33.
8. Saito T, Shimazaki Y, Koga T, Tsuzuki M and Ohshima A: Relationship between periodontitis and hepatic condition in Japanese women. *J Int Acad Periodontol* 2006;8:89-95.
9. Morita T, Yamazaki Y, Fujiharu C, Ishii T, Seto M, Nishinoue N et al, Serum γ -glutamyltransferase level is associated with periodontal disease independent of drinking habits in Japanese adults. *Med Sci Monit* 2014;20:2109-16.
10. Kumar PS, Griffen AL, Barton JA, Paster BJ, Moeschberger ML and Leys EJ et al. New bacterial species associated with chronic periodontitis. *J Dent Res* 2003;82:338-44.
11. Kumar PS, Griffen AL, Moeschberger ML and Leys EJ: Identification of candidate periodontal pathogens and

- beneficial species by quantitative 16S clonal analysis. *J Clin Microbiol* 2005;43:3944-55.
12. Fernández J and Gustot T: Management of bacterial infections in cirrhosis. *J Hepatol* 2012;56(1):S1-S12.
 13. and Rodés J: Bacterial infections in cirrhosis: Epidemiological changes with invasive procedures and norfloxacin prophylaxis. *Hepatol* 2002;35:140-48.
 14. Boggess KA, Madianos PN, Preisser JS, Moise KJ Jr and Offenbacher S: Chronic maternal and fetal *Porphyromonas gingivalis* exposure during pregnancy in rabbits. *Am J Obstet Gynecol* 2005;192:554-7.
 15. Raja M, Ummer F and Dhivakar CP: *Aggregatibacter actinomycetemcomitans* - a tooth killer? *J Clin Diagn Res* 2014;8:ZE13-ZE16.
 16. Narayanan SK, Nagaraja TG, Chengappa MM and Stewart GC: Leukotoxins of gram-negative bacteria. *Vet Microbiol* 2002;84:337-56.
 17. Chen PB, Davern LB, Neiders ME, Reynolds HS and Zambon JJ: Analysis of in vitro lymphoproliferative responses and antibody formation following subcutaneous injection of *Actinobacillus actinomycetemcomitans* and *Wolinella recta* in a murine model. *Oral Microbiol Immunol* 1991;6:12-6.
 18. Tuomainen AM, Jauhiainen M, Kovanen PT, Metso J, Paju S and Pussinen PJ et al. *Aggregatibacter actinomycetemcomitans* induces MMP-9 expression and proatherogenic lipoprotein profile in apoE-deficient mice. *Microb Pathog* 2008;44:111-7.
 19. Lentsch AB, Yoshidome H, Kato A, Warner RL, Cheadle WG, Ward PA et al. Requirement for interleukin-12 in the pathogenesis of warm hepatic ischemia/reperfusion injury in mice. *Hepatol* 1999;30:1448-53.
 20. Gandoura S, Weiss E, Rautou PE, Fasseu M, Gustot T, Lemoine F et al. Gene- and exon-expression profiling reveals an extensive LPS-induced response in immune cells in patients with cirrhosis. *J Hepatol* 2013;58:936-48.
 21. Moreau R, Barrière E, Tazi KA, Lardeux B, Dargère D, Urbanowicz W et al, Terlipressin inhibits in vivo aortic iNOS expression induced by lipopolysaccharide in rats with biliary cirrhosis. *Hepatol* 2002;36:1070-8.
 22. Abu-Shanab A and Quigley EM: The role of the gut microbiota in nonalcoholic fatty liver disease. *Nat Rev Gastroenterol Hepatol* 2010;7:691-701. Madianos PN, Bobetsis YA
 23. Madianos PN, Bobetsis YA and Kinane DF: Generation of inflammatory stimuli: How bacteria set up inflammatory responses in the gingiva. *J Clin Periodontol* 2005;32(6):57-71.
 24. Malik R, Mookerjee RP and Jalan R: Infection and inflammation in liver failure: Two sides of the same coin. *J Hepatol* 2009;51:426-9.
 25. Muñoz NM, Katz LH, Shina JH, Gi YJ, Menon VK, Gagea M et al, Generation of a mouse model of T-cell lymphoma based on chronic LPS challenge and TGF- β signaling disruption. *Genes Cancer* 2014;5:348-52.
 26. Su GL, Klein RD, Aminlari A, Zhang HY, Steinstraesser L, Alarcon WH et al, Kupffer cell activation by lipopolysaccharide in rats: Role for lipopolysaccharide binding protein and toll-like receptor 4.
 27. Medzhitov R: Toll-like receptors and innate immunity. *Nat Rev Immunol* 2001;1:135-45.
 28. Mattsson E, Verhage L, Rollof J, Fleer A, Verhoef J and van Dijk H et al. Peptidoglycan and teichoic acid from *Staphylococcus epidermidis* stimulate human monocytes to release tumour necrosis factor-alpha, interleukin-1 beta and interleukin-6. *FEMS Immunol Med Microbiol* 1993;7:281-7.
 29. Forner L, Nielsen CH, Bendtzen K, Larsen T and Holmstrup P: Increased plasma levels of IL-6 in bacteremic periodontitis patients after scaling. *J Clin Periodontol* 2006;33:724-9.
 30. Loos BG, Craandijk J, Hoek FJ, Wertheim-van Dillen PM and van der Velden U: Elevation of systemic markers related to cardiovascular diseases in the peripheral blood of periodontitis patients. *J Periodontol* 2000;71:1528-34.

How to cite this article: Hassan SA. Periodontitis and liver disease a relation – Review. *J Urol Nephrol Hepatol Sci* 2019;2(4):53-6.