

## Sanguinaria-a plant extract is safe alternative than Triclosan - As an antiplaque agent

Sauvik Singha<sup>1</sup>, Jugajyoti Pathi<sup>2\*</sup>, Anshalika Agrawal<sup>3</sup>, Jay Taank<sup>4</sup>, Irfan Zunzani<sup>5</sup>, Kshitij Aggarwal<sup>6</sup>

<sup>1,3-6</sup>PG Resident <sup>2</sup>Senior Lecturer, <sup>1-6</sup>Dept. of Oral and Maxillofacial Surgery, <sup>1,3-6</sup>Hitkarini Dental College and Hospital, Jabalpur, Madhya Pradesh, <sup>2</sup>Kalinga Institute of Dental Sciences, Bhubaneswar, Odisha, India

\*Corresponding Author: Jugajyoti Pathi

Email: jpathi@kims.ac.in

### Abstract

Triclosan is a polychloro phenoxy phenol., an antiplaque agent used in toothpaste and mouth rinse (Colgate Total, Janina Diamond Whitening Toothpaste, Breeze TM Triclosan Mouthwash, and Reach Antimicrobial Toothbrush). But it has questionable benefits in toothpastes and the negative effects on endocrine system, immune system, musculoskeletal system and environment. Also causes allergy and carcinogenesis. In context to that Triclosan, Sanguinarine is a safe alternative (mouthwash Viadent (USA) Perioguard (UK) Sanguinarine, alkaloids, main a bioactive component of bloodroot, *Sanguinaria Canadensis* L., is a member of Papaveraceae family. It has antiseptic, anti-inflammatory and antiplaque agent. Part of its appeal would be that it is a *natural therapeutic product*, as distinct from a 'synthetic' or 'artificial' chemical.

**Keywords:** Sanguinaria, Triclosan, Bloodroot, Gingivitis, Plaque.

### Introduction

The Prevalence of gingivitis in adults in the UNITED STATES exceeds 50% and approaches 100% in some population group.<sup>1</sup> Plaque considered the key factor contributing to gingival inflammation that, if left untreated may progress to periodontitis.

Triclosan is used as antiplaque agent in toothpaste as well as mouth rinse [Colgate Total, Janina Diamond Whitening Toothpaste, Breeze TM Triclosan Mouthwash, Reach Antimicrobial Toothbrush]. Chemically it is a polychloro phenoxy phenol.<sup>2</sup> The negative effects of Triclosan on the environment and its questionable benefits in toothpastes<sup>3</sup> have led to the Swedish Naturskyddsföreningen to recommend not using Triclosan in toothpaste.<sup>4</sup>




Fig. 1:

### Acute Effects

Triclosan has also been associated with a higher risk of food allergy.<sup>5</sup> Other studies have linked Triclosan to allergic contact dermatitis in some individuals.<sup>6</sup> The American Dental Association published a response to the concerns stemming from the Virginia Tech study stating that the study is not relevant to toothpaste.<sup>7</sup> Manufacturers of a number of Triclosan containing toothpaste claims that it active ingredient continues to work for up to 12 hours after its use.

### Chronic Effects

#### On Endocrine

A 2006 study concluded that low doses of Triclosan act as an endocrine disruptor in the North American bullfrog.<sup>8</sup> The hypothesis proposed that Triclosan blocks the metabolism of thyroid hormone because it chemically mimics thyroid hormone. A study between 2003 and 2006 concluded that Triclosan affects the immune system and showed a positive association with allergy or hay fever diagnosis.<sup>9</sup> Another 2009 study demonstrated that Triclosan exposure significantly impacts thyroid hormone concentrations in the male juvenile rats.<sup>10</sup> Triclosan is also showing up in dolphins near South Carolina and Florida in concentrations known to disrupt hormones, growth, and development in other animals.<sup>11</sup> *On Muscle contraction*-It impairs excitation-contraction coupling in cardiac and skeletal muscle function in mice.<sup>12</sup> *carcinogenesis*-Triclosan can combine with chlorine to form chloroform,<sup>13</sup> which the United States Environmental Protection Agency classifies as a probable human carcinogen.

#### Link to Dioxin

Dioxin is highly carcinogenic and cause weakening of immune system, altered sex hormones, miscarriage, cancer.<sup>[14]</sup> Triclosan is listed as 'could be' and 'suspected to be' contaminated with dioxin in EPA's draft Dioxin Reassessment.

#### Resistance

*Escherichia coli* and *Staphylococcus aureus* develop low-level resistance to Triclosan at its lower bacteriostatic concentrations because of *FabI* mutations, which results in a decrease of triclosan's effect on ENR-NAD<sup>+</sup> binding<sup>15,16</sup> It also gain low-level resistance to Triclosan is to over express *FabI*.<sup>17</sup> *Pseudomonas aeruginosa* have innate resistance to Triclosan at low, bacteriostatic levels,<sup>18</sup> *Bacillus* genus, have alternative *FabI* genes

(FabK) to which Triclosan does not bind and hence are less susceptible.



Fig. 2:

### Environmental Effects

Triclosan effluents affect both the structure and the function of algal communities in stream ecosystems. Because of its lipophilic nature and resistance to degradation, Triclosan in waterways is readily available for absorption, bioaccumulation by aquatic organisms in environment.<sup>19</sup>

*In context to that Triclosan is a safe alternative is Sanguinarine. Part of its appeal would be that it is a natural therapeutic product, as distinct from a 'synthetic' or 'artificial' chemical.*

Sanguinarine has antiseptic, anti-inflammatory and antiplaque agent.<sup>20</sup> Sanguinarine is a benzophenanthridine alkaloid derived from rhizomes of *Sanguinaria Canadensis* L. It is a cationic molecule which converts from an iminium ion form at pH less than 6 to an alkanolamine form at pH greater than 7. *Sanguinaria* extract is composed of Sanguinarine and five other closely related alkaloids. The safety profile of both Sanguinarine and *sanguinaria* extract provide a broad margin for their safe use in oral health products. Sanguinarine has broad antimicrobial activity as well as anti-inflammatory properties.<sup>21</sup> It has been incorporated into dentifrices and mouthwash Viadent (USA) Perioguard (UK). Bloodroot is a member of Papaveraceae family. It is a native spring wildflower that grows in rich wood lands of North America

### Uses

Currently it is being studied for use as an anticancer agent. Mainly it has commercial success in toothpaste and mouthwash as an anti-plaque agent. The United States FDA has approved the inclusion of Sanguinarine in toothpaste as an antibacterial or antiplaque agent.<sup>22,23</sup>

In a short-term double-blind cross-over study of plaque re-growth and gingivitis lasting two weeks for each experimental period, using a mouth rinse which was a 0.03% solution of *sanguinaria* extract. They controlled the study with a placebo with similar taste and color. Plaque indices were 40% lower and gingivitis scored 25% lower with the *sanguinaria* rinses compared with the controls. Normal tooth brushing was suspended during the trial period, and with some of the subjects there was a slight

discoloration of the teeth and dorsum of the tongue. [Wennstrom and Lindhe (1985)]

In a six-month double-blind parallel study, assessed gingivitis and plaque in subjects using Viadent dentifrice together with a mouth rinse with *sanguinaria* extract and zinc chloride. The negative controls used the same products but without the *sanguinaria* extract or zinc chloride. Plaque and gingivitis scores in the *sanguinaria* extract group were 21% and 25% lower than those in the control group at the end of 6 months. "Bleeding on probing" assessments also showed a reduction in the test group compared with controls. No opportunistic overgrowth of pathogens was found, but there were reductions in numbers of the organisms associated with gingivitis, which may have accounted for the reduction in gingivitis. [Harper *et al.* (1990a)(1990b)]

This report represents the findings of an Expert Panel on the safety of *Sanguinaria* extract used in Viadent oral rinse and toothpaste products and represents an independent review of the *Sanguinaria* extract toxicological data base. The Panel concluded that the data base on *Sanguinaria* extract is substantial and indicates that *Sanguinaria* extract is safe in its present use in Viadent products based on a large margin of safety between levels of human exposure and levels found to produce minimum effect or to be without adverse effect in animals.<sup>24</sup>

Short- and long-term testing of *sanguinaria* toothpaste and oral rinse used individually have yielded both positive and negative results. This review evaluates the results of a number of clinical trials testing the regimen use of *sanguinaria* products for periods ranging from 14 days to six months. Review of these trials establishes the clinical efficacy of the two products in combination. The regimen produces positive reductions in plaque, gingival inflammation and bleeding parameters for up to six months with no adverse hard tissue effects and only one reversible adverse soft tissue effect observed among the 260 subjects tested. In addition, no adverse microbiological shifts in the normal oral flora were observed.<sup>25</sup>

In vitro studies indicate that the anti-plaque action of *sanguinaria* is due to its ability to inhibit bacterial adherence to newly formed pellicle, its retention in plaque being 10-100 times its saliva concentration, and due to its antimicrobial properties. The MIC of Sanguinarine ranges from 1 to 32 micrograms/mL for most species of plaque bacteria. Long term use of *sanguinaria*-containing toothpaste and oral rinse products does not predispose users to detrimental shifts in oral flora. Electron microscopic studies of bacteria exposed to Sanguinarine demonstrate that bacteria aggregate and become morphologically irregular. Sanguinarine-containing slow release polymer systems are currently being developed for use in periodontitis treatment applications.<sup>20</sup>

### Acknowledgment

The authors would like to thank Dr. Roberto Vienna, Past FDI President, Dr Viney Aggarwal, Consultant Oral and Maxillofacial Surgeon, Aggarwal Dental Clinic.

**Conflict of Interest:** None.

## Reference

- American Medical Association. 2000. Use of antimicrobials in consumer products. Report 2 of the Council on Scientific Affairs(A-00)
- Triclosan safety is currently under review by the FDA, and Health Canada.
- Edvardsson S, Burman LG, Adolfsson-Erici M, Bäckman N. "Risker och nytta med triklosan i tandkräm" [Risks and benefits of triclosan in toothpaste]. *Tandläkartidningen* 97(10):58–64.
- Start ~ Naturskyddsforeningen
- Sicherer, SH; Leung, DY. "Advances in allergic skin disease, anaphylaxis, and hypersensitivity reactions to foods, drugs, and insects in 2012". *J Allergy Clin Immunol* 2013;131(1):55–66. doi:10.1016/j.jaci.2012.11.007. PMID 23199604
- Bhutani T, Jacob SE. "Triclosan: a potential allergen in suture-line allergic contact dermatitis". *Dermatol Surg* 2009;35(5):888–9. doi:10.1111/j.1524-4725.2009.01151.x.PMID 19389086.
- Yang, Jeniffer. "Experts concerned about dangers of antibacterial products". *Globe and Mail*. Retrieved 2009-08-25.
- Veldhoen N, Skirrow RC, Osachoff H. "The bactericidal agent triclosan modulates thyroid hormone-associated gene expression and disrupts postembryonic anuran development". *Aquat Toxicol* 2006;80(3):217–27. doi:10.1016/j.aquatox.2006.08.010.PMID 17011055.
- Clayton EM, Todd M, Dowd JB, Aiello AE (March). "The impact of bisphenol A and triclosan on immune parameters in the U.S. population, NHANES 2003-2006". *Environ. Health Perspect* 2011;119(3):390–6. doi:10.1289/ehp.1002883. PMC 3060004.PMID 21062687.
- Zorrilla LM, Gibson EK, Jeffay SC. "The effects of triclosan on puberty and thyroid hormones in male Wistar rats". *Toxicol Sci* 2009;107(1):56–64. doi:10.1093/toxsci/kfn225. PMID 18940961.
- "Germ-killing chemical from soaps, toothpaste building up in dolphins". 2009-08-11. Retrieved 2011-04-11.
- Cherednichenko G, Zhang R, Bannister RA. "Triclosan impairs excitation-contraction coupling and Ca<sup>2+</sup> dynamics in striated muscle". *Proc Natl Acad Sci U.S.A.* 2012;109 (35):14158–63. Bibcode:2012PNAS..10914158C.doi:10.1073/pnas.1211314109. PMC 3435154. PMID 22891308
- Yang, Jeniffer. "Experts concerned about dangers of antibacterial products". *Globe and Mail*. Retrieved 2009-08-25.
- US Dept of health and human Service 1988 Toxicological profile for chlorinated dibenzo-p-dioxins.
- Heath RJ, Rubin JR, Holland DR, Zhang E, Snow ME, Rock CO. "Mechanism of triclosan inhibition of bacterial fatty acid synthesis". *J Biol Chem* 1999;274(16):11110–4. doi:10.1074/jbc.274.16.11110. PMID 10196195.
- Fan F, Yan K, Wallis NG. "Defining and combating the mechanisms of triclosan resistance in clinical isolates of *Staphylococcus aureus*". *Antimicrob. Agents Chemother* 2002;46(11):3343–7. doi:10.1128/AAC.46.11.3343-3347.2002. PMC 128739.PMID 12384334.
- Slater-Radosti C, Van Aller G, Greenwood R. "Biochemical and genetic characterization of the action of triclosan on *Staphylococcus aureus*". *J Antimicrob Chemother* 2001;48 (1): 1–6. doi:10.1093/jac/48.1.1. PMID 11418506.
- Chuanchuen R, Karkhoff-Schweizer RR, Schweizer HP. "High-level triclosan resistance in *Pseudomonas aeruginosa* is solely a result of efflux". *Am J Infect Control* 2003;31(2):124–7. doi:10.1067/mic.2003.11. PMID 12665747.
- Wilson BA, V H Smith, F deNoyelles jr CK Larvie. Effects of three pharmaceutical and personal care product on natural fresh water algal assemblages. *Environ Sci Technol* 2003;37(9):162A-164A
- Cech R. 2002 Growing At risk Medicinal Plants. Horizon herbs. Williams
- Antimicrobial action of sanguinarine. Godowski KC. *J Clin Dent* 1989;1(4):96-101.
- Godowski, KC. "Antimicrobial action of sanguinarine". *J Clin Dent* 1989;1(4):96–101. PMID 2700895.
- Southard, GL, Boulware, RT, Walborn, DR; Groznik, WJ; Thorne, EE; Yankell, SL. "Sanguinarine, a new antiplaque agent: Retention and plaque specificity". *J Am Dent Assoc* 1984;108(3):338–41. PMID 6585404.
- Safety of Sanguinaria extract as used in commercial toothpaste and oral rinse products. Frankos VH, Brusick DJ, Johnson EM, Maibach HI, Munro I, Squire RA, Weil CS. *J Can Dent Assoc* 1990;56(7):41-7.
- Kuftinec MM, Mueller-Joseph LJ, Kopczyk RA. Sanguinaria toothpaste and oral rinse regimen clinical efficacy in short- and long-term trials. *J Can Dent Assoc* 1990;56(7):31-3.

**How to cite this article:** Singha S, Pathi J, Agrawal A, Taank J, Zunzani I, Aggarwal K. Sanguinaria-a plant extract is safe alternative than Triclosan - As an antiplaque agent. *Int J Aesthet Health Rejuvenation* 2019;2(2):20-22.